


*Osteopathic management of knee joint symptoms – a brief summary of current literature*

Osteopathic care contains over 100 different techniques or procedures<sup>1,2,3,4,5</sup>. It is notable in treatment of knee joint symptoms that a wide range of techniques are used.



**Lateral view of the knee joint.**

Image supplied by the National Institute of Arthritis, Musculoskeletal and Skin Diseases.  
[www.niam.nih.gov](http://www.niam.nih.gov).

The most commonly used techniques are broadly grouped into seven major types:

- High velocity low amplitude (HVLA)
- Soft tissue/massage techniques<sup>6</sup>
- Articulation involving gentle repetitive movement of the joint.
- Muscle energy involving repeated isometric contractions with passive joint

movement<sup>2,3</sup>.

- Counterstrain involving the symptomatic joint being placed in a position of least discomfort while at the same time monitoring the degree of tenderness at a nearby tender point until the tenderness reduces<sup>2,3,7</sup>.
- Myofascial release techniques to stretch muscle and reduce tension<sup>3</sup>.
- Lymphatic pump techniques to mechanically assist lymphatic drainage<sup>8</sup>.

Studies focussing on the osteopathic management of the knee joint describe the use of a range of techniques. The number of published studies is much smaller than for other areas of the body. The studies undertaken include a range involving animal studies, laboratory work, intervention studies, case studies and opinion pieces/commentaries, and randomised controlled trials.

Animal studies have focussed mainly on the knee joint in the rabbit. Laboratory studies include investigation of the effects of techniques to correct measurements of the knee angle, the physiological effects of the lack of weight on the knee, the introduction of surfactants in the early management of cell death of the articular cartilage following knee trauma, and the effect of fatigue on different muscle groups (notably the hamstrings).

Most of the published studies on knee symptoms have been undertaken in American osteopathic institutions and have investigated the effect of osteopathic management on patients after surgery. These studies have looked at the effects of different drug management programmes, and different surgical approaches in terms of fixation media and the usefulness of unilateral or bilateral knee arthroplasty. One study looked at the use of osteopathic manipulative treatment in patients following either hip or knee arthroplasty<sup>9</sup>. The protocol used in the study involved myofascial release, strain/counterstrain, muscle energy, soft tissue, high velocity low amplitude (HVLA) manipulation, and craniosacral techniques. The treatments administered in the study were not found to be efficacious for the hospital-based patient population.

A small number of published clinical studies were identified; they are summarised in the table overleaf.



Title of study	Type of study	Intervention	Population size (N)	Summary of findings	Link
Brantingham <i>et al.</i> Manipulative therapy for lower extremity conditions <sup>10</sup>	Literature review	Manual therapy using a variety of approaches	N/A	A total of 39 relevant peer-reviewed papers were identified for this literature review demonstrating management of a number of lower extremity disorders.	<a href="http://www.ncbi.nlm.nih.gov/pubmed/19121464">http://www.ncbi.nlm.nih.gov/pubmed/19121464</a>
Opila-Correia KA. Kinematics of high-heeled gait <sup>11</sup>	Investigational study	Measurement of three-dimensional kinematics of the tibia, knee, hip, pelvis, trunk and upper trunk for high or low-heeled gait	N=14	High-heeled gait subjects had shorter stride lengths, walked more slowly, and had higher stance time. Knee flexion was increased at heel strike and during stance phase; lower knee and hip flexion occurred during swing phase, and lower range of motion of the pelvis in the sagittal plane,	<a href="http://www.ncbi.nlm.nih.gov/pubmed/2327881">http://www.ncbi.nlm.nih.gov/pubmed/2327881</a>
Campos <i>et al.</i> Muscular adaptations in response to three different resistance-	Investigational study	An eight week progressive resistance training programme to investigate the	N=32	Maximal aerobic power increased for HR group only, all three major muscle fibre types hypertrophied for the LR and IR groups, and the HR group	<a href="http://www.ncbi.nlm.nih.gov/pubmed/12436270">http://www.ncbi.nlm.nih.gov/pubmed/12436270</a>

training regimens <sup>12</sup>		<p>strength-endurance continuum. Interventions included low repetitions (LR)[3-5 sets of reps with 3 mins rest between 4 sets], intermediate repetitions (IR) [9-11 reps, 2 mins rest between 3 sets], high repetition (HR)[20-28 reps with 1 min rest between 2 sets), and a non-exercising Control (CTL).</p>		<p>adapted better for submaximal prolonged contractions.</p>	
<p>Meyer <i>et al.</i> Excessive compression of the human tibio-femoral joint causes ACL rupture<sup>13</sup></p>	<p>Laboratory study on previously asymptomatic knees from fresh cadavers.</p>	<p>Excessive axial compression load was applied to the tibio-femoral joint at 60, 90 or 120° of flexion</p>	<p>N=16</p>	<p>The maximum force for ACL failure was 5.1kN for all flexion angles combined; at 90° flexion injury occurred with relative anterior displacement, lateral displacement and internal rotation of the tibia on the femur.</p>	<p><a href="http://www.ncbi.nlm.nih.gov/pubmed/16154419">http://www.ncbi.nlm.nih.gov/pubmed/16154419</a></p>

McClinton <i>et al.</i> Influence of step height on the quadriceps onset timing and activation during stair descent in individuals with patellofemoral pain syndrome (PFPS) <sup>14</sup>	Case control study	Data concerning knee kinematics and quadriceps activity was collected during ascending of 5 different step heights	N=20	Quadriceps onset timing and magnitude was similar regardless of step height between subjects with or without (PFPS)	<a href="http://www.ncbi.nlm.nih.gov/pubmed/17549952">http://www.ncbi.nlm.nih.gov/pubmed/17549952</a>
Meyer and Haut. ACL injury induced by internal tibial torsion or tibiofemoral compression <sup>15</sup>	Laboratory study involving knees from 7 cadavers	Compression or torsion experiments were conducted to assess if the magnitude and type of motion before ACL rupture would significantly change from just before ACL rupture.	N=7	ACL injury was documented in all knees at 5.4kN of compression or 33Nm of internal tibial torque.	<a href="http://www.ncbi.nlm.nih.gov/pubmed/19007932">http://www.ncbi.nlm.nih.gov/pubmed/19007932</a>
Jarski <i>et al.</i> The effectiveness of OMT treatment as complementary therapy following	Prospective match-controlled outcome study	Treatment group received OMT on postoperative days 2 -5.	N=76	Participants receiving osteopathic care in the early postoperative period negotiated stairs earlier and walked greater distances than the control group	<a href="http://www.ncbi.nlm.nih.gov/pubmed/10979164">http://www.ncbi.nlm.nih.gov/pubmed/10979164</a>

surgery <sup>16</sup>				participants	
Smith and Fryer. A comparison of two muscle energy techniques for increasing flexibility of the hamstring muscle group <sup>17</sup>	Laboratory based investigational study	Muscle energy technique (MET) was applied with 30 s post-isometric stretch phase, or with 3s post-isometric stretch phase. Hamstring measurement was undertaken using active knee extension (AKE).	N=40	Both techniques appeared to be equally effective in increasing hamstring extensibility.	<a href="http://www.ncbi.nlm.nih.gov/pubmed/19083689">http://www.ncbi.nlm.nih.gov/pubmed/19083689</a>
Barron and Rubin. Managing osteoarthritic knee pain <sup>18</sup>	Commentary		N/A	Pharmacological and non-pharmacological approaches to treating patients with osteoarthritic pain are discussed.	<a href="http://www.ncbi.nlm.nih.gov/pubmed/17986674">http://www.ncbi.nlm.nih.gov/pubmed/17986674</a>
Gugel and Johnston. Osteopathic manipulative treatment of a 27 year old man after ACL reconstruction <sup>19</sup>	Case study	Somatic dysfunction was identified in the lumbopelvic region, in addition to increased muscular tension around the injured knee, ankle	N=1	OMT was used post surgery. Increased mobility in the lumbopelvic region was recorded, areas of somatic dysfunction resolved and the patient was able to return to regular sporting activity 6	<a href="http://www.ncbi.nlm.nih.gov/pubmed/16790541">http://www.ncbi.nlm.nih.gov/pubmed/16790541</a>

		joint, lower thorax and between ribs 6-9.		months after surgery.	
Rubin. Management of osteoarthritic knee pain <sup>20</sup>	Commentary		N/A	Pharmacological and non-pharmacological approaches to treating patients with osteoarthritic pain are discussed.	<a href="http://www.ncbi.nlm.nih.gov/pubmed/16249363">http://www.ncbi.nlm.nih.gov/pubmed/16249363</a>
Meyer <i>et al.</i> Osteochondral microdamage from valgus bending of the human knee <sup>21</sup>	Laboratory based investigational study	Four pairs of knees were loaded in valgus bending until gross injury occurred. Peak valgus movement and resultant movement of the joint were recorded. Pressure sensitive film documented the location and magnitude of tibiofemoral contact. Micro-CT scans identified microcracks in the	N=8 (knees)	Peak bending with ligamentous failure occurred at 107Nm. Cartilage fissures and subchondral bone microcracks occurred in areas of high contact pressure.	<a href="http://www.ncbi.nlm.nih.gov/pubmed/19505750">http://www.ncbi.nlm.nih.gov/pubmed/19505750</a>

		subchondral bone.			
Cheng and Shi. Rehabilitation exercises after single total knee replacement <sup>22</sup>	Case series	Postoperative rehabilitation raining was undertaken at a Chinese osteopathic hospital	N=38	The authors recorded that the rehabilitation programme had produced satisfactory results and recommended wider implementation.	<a href="http://www.ncbi.nlm.nih.gov/pubmed/20415085">http://www.ncbi.nlm.nih.gov/pubmed/20415085</a>
Pedowitz RN. Use of osteopathic manipulative treatment for iliotibial band friction syndrome <sup>23</sup>	Case study	Treatment used osteopathic techniques but specifically counterstrain technique	N=1	A tender point was identified 0-3cm proximal to the lateral femoral condyle and was treated accordingly.	<a href="http://www.jaoa.org/cgi/content/full/105/12/563">http://www.jaoa.org/cgi/content/full/105/12/563</a>

There are a large number of laboratory and post-surgical trials published. There is a notable absence of published work documenting the various management techniques used in everyday osteopathic practice for our patients. The National Council for Osteopathic Research will be undertaking a data collection project in 2011 to try and address this paucity of data. An email invitation to participate will be circulated to osteopaths.

## References

1. Lesho, E. P. An overview of osteopathic medicine. *Arch Fam Med.* 1999;8,477-84.
2. DiGiovanna EL, Martinke DJ, Dowling DJ. Introduction to osteopathic medicine. In DiGiovanna EL, Sciowitz S eds. *An Osteopathic Approach to Diagnosis and Treatment.* Philadelphia Pa: JB Lippincott;1991:1-31.
3. Greenman PE. *Principles of Manual Medicine.* Baltimore, Md: Williams and Wilkins; 1989:1-13,30.
4. Still AT. *Osteopathy Research and Practice.* Seattle, Walsh: Eastland Press;1992:xxi-13.
5. Owens C. *Endocrine Interpretation of Chapman's Reflexes.* Newark, Ohio: American Academy of Osteopathy; 1963.
6. Furlan AD, Imamura M, Dryden T *et al.* Massage for low back pain: an updated systematic review within the framework of the Cochrane Back Review Group. *Spine.* 2009;34(16):1669-84.
7. Jones HL. *Strain and Counterstrain.* Newark, Ohio: American Academy of Osteopathy; 1981.
8. Degenhardt BF, Kuchera JL. Update on osteopathic medicine concepts and the lymphatic system. *J Am Osteopath Assoc.* 1996;96:97-100.
9. Licciardone JC, Stoll ST, Cardarelli KM *et al.* A randomised controlled trial of osteopathic manipulative treatment following knee or hip arthroplasty. *Journal of the American Osteopathic Association.* 2004;104(5):193-202.
10. Brantingham JW, Globe G, Pollard H *et al.* Manipulative Therapy for lower extremity conditions: expansion of literature review. *Journal of Manipulative and Physiological Therapeutics.* 2009;32:53-7.
11. Opila-Correia KA. Kinematics of high heeled gait. *Archives of Physical Medicine and Rehabilitation.* 1990;71(5):304-9
12. Campos GE, Luecke TJ, Wendeln HK *et al.* Muscular adaptations in response to three different resistance-training regimens: specificity of repetition maximum training zones. *European Journal of Applied Physiology.* 2002;88(1-2):50-60

13. Meyer EG, Haut RC. Excessive compression of the human tibio-femoral joint causes ACL rupture. *Journal of Biomechanics*. 2005;38:2311-2316.
14. McClinton S, Donatell G, Weir J *et al*. Influence of step height on quadriceps onset timing and activation during stair ascent in individuals with patellofemoral pain syndrome. *Journal of Orthopaedic and Sports Physical Therapy*. 2007;37(5):239-44.
15. Meyer EG, Haut RC. Anterior cruciate ligament injury induced by internal tibial torsion or tibiofemoral compression. *Journal of Biomechanics*. 2008;41:3377-3383
16. Jarski RW, Loniewski EG, Williams J *et al*. The effectiveness of osteopathic manipulative treatment as complementary therapy following surgery: a prospective, match-controlled outcome study. *Alternative Therapy Health and Medicine*. 2000;6(5):77-81.
17. Smith M, Fryer G. A comparison of two muscle energy techniques for increasing flexibility of the hamstring muscle group. *Journal of Bodywork and Movement Therapies*. 2008;12(4):312-7.
18. Barron MC, Rubin BR. Managing osteoarthritic knee pain. *Journal of the American Osteopathic Association*. 2007;107(11):ES21-ES27.
19. Gugel MR, Johnston WL. Osteopathic manipulative treatment of a 27 year old man after anterior cruciate ligament reconstruction. *Journal of the American Osteopathic Association*. 2006;106(6):346-349.
20. Rubin BR. Management of osteoarthritic knee pain. *Journal of the American Osteopathic Association*. 2005;105(9):S23-S28.
21. Meyer EG, Villwock MR, Haut RC. Osteochondral microdamage from valgus bending of the human knee. *Clinical Biomechanics*. 2009;24(7):577-82.
22. Cheng N, Shi QY. Rehabilitation exercises after single total knee replacement: a report of 38 cases. *Zhongguo Gu Shang*. 2010;23(3):220-1.
23. Pedowitz RN. Use of osteopathic manipulative treatment for iliotibial band friction syndrome. *Journal of the American Osteopathic Association*. 2005;105(12):5