

The use of ice in patient management – a brief review the evidence

Carol Fawkes, NCOR Research Officer

The use of cold, or cryotherapy, for medicinal purposes in the form of ice and snow has been used since the time of Hippocrates¹. And this treatment strategy is still often used in osteopathic practice, as well as many other healthcare disciplines. The application of ice in osteopathic treatment is most commonly recommended for patients presenting with musculoskeletal injury^{2,3}: ice is, however, frequently used post-operatively⁴, in rheumatic disease⁵ and occasionally for haemophilia⁶ patients.

Proposed physiological processes

The rationale for using ice is to achieve a number of outcomes; these commonly centre around the processes that occur during an acute inflammatory process:

- *Reduction of oedema to reduce compartmental pressure.* Theoretically, the application of cold should cause vasoconstriction in superficial blood vessels, contributing to the reduction in oedema^{7,8}. This point of view has, however, been challenged by Curl *et al*⁹ who suggested an alternative mechanism must be operating. Cold is also believed to have a secondary nociceptive effect by limiting oedema formation and thereby limiting painful tissue distention.
- *Reduction of haemorrhage and haematoma formation.* Ho¹⁰ and colleagues identified that applying ice for 20 minutes significantly decreased arterial and soft tissue blood flow around the knee joint; bone blood flow and metabolism were also significantly reduced. Paradoxically, periods of vasoconstriction are found to be followed by vasodilation and then vasoconstriction again. Grana *et al*¹¹ suggest that vasoconstriction would occur until subcutaneous temperatures fall below 15 degrees Celsius; this would then be followed by vasodilation, caused either by paralysis of the contractile mechanism or blockage of the constrictor signals. This dilation process is thought to account for the reddening that occurs when cold is applied to the skin surface.
- *Hypoxic tissue damage is reduced as the temperature reduction slows the metabolic rate of the injured tissue and reduces the rate of oxygen use¹².*
- *Muscle spasm is reduced due to the cold exerting an inhibitory effect on the muscle spindles.*
- *Pain is diminished due to a combination of three different processes:*
 - Cold-induced neurapraxia, where sensory nerve impulses are completely blocked or slowed down.
 - The cycle of pain and muscle spasm is interrupted as the reduction of muscle spasm diminishes levels of pain.
 - Loeser¹³ suggested that the application of cold causes signals to be transmitted to the spinal cord, which override pain impulses as they enter the spinal cord.

Reviewing the evidence

Chronic pain is less commonly associated with the use of cold packs, however, the pain-reducing effect of cold application can be helpful in some chronic pain

conditions¹⁴. Healthcare practitioners frequently advocate the use of ice in acute symptom management post-surgery and particularly so in the care of sports injuries. One study comparing the use of ice with no ice in patients recovering from arthroscopic knee surgery found that patients using ice had significantly lower pain scores, when measured using the McGill Pain Questionnaire, and used significantly less prescription and non-prescription analgesia¹⁵.

Reviews have been carried out to investigate whether cold therapy does hasten a return to sporting activity. However, the reviewers were concerned at the quality of the studies addressing this therapeutic area; they concluded that cryotherapy may have a positive effect on return to sport participation, but suggested that considerably more work was required to raise the quality of future studies and a greater need was identified for the use of return-to-participation outcome measures¹⁶. Various studies have looked at specific injuries associated with sportsmen and women and the use of ice in their management is a common thread¹⁷.

Compression

There is little evidence to suggest that the combination of ice and compression has any significant effect. Studies are predominantly limited to the treatment of hospital inpatients and very few studies have assessed the management of closed soft-tissue injuries. The studies involving ice and compression fail to use the same modes of compression, duration of ice and mode of ice application. It is therefore impossible to draw any objective information from the studies that add value to clinical practice¹⁸.

Type of ice pack

Ice packs have become increasingly sophisticated; some patients still prefer to use good old-fashioned frozen peas, while others find that flexible gel packs are more convenient, especially when travelling while experiencing pain. Chesterton *et al*¹⁹ compared the localised skin-cooling effects of an ice pack made from frozen peas with a flexible frozen gel pack. They found that the latter failed to cool the skin as well as the frozen peas and did not produce an effect that induced localised skin analgesia, reduced nerve conduction velocity and reduced metabolic enzyme levels to clinically relevant levels.

Zemke *et al*²⁰ compared the effect of ice pack application with ice massage. Ice massage involved an ice cup being massaged over the gastrocnemius muscle using overlapping horizontal strokes for 15 minutes. Ice massage reached a lower temperature of 17.9°C in 2.4 minutes and the ice pack achieved a temperature of 28.2°C in 12.5 minutes.

When to apply ice

In the 1940's²¹ the recommendation was to apply ice within the first 30-60 minutes after injury. By the 1950's this advice had changed to within the first 24-72 hours after injury²².

Duration of application

Stitik and Nadler²³ recommend application of ice for 20-30 minutes every two hours, combined with rest and compression; they recommend this treatment to be

continued until the swelling has gone down or after 48 hours has passed. They propose that the “20 minutes on, 20 minutes off” regime potentially increases the risk of thermal injury.

Hochberg⁴ compared the effect of continuous ice application with that of intermittent application for patients following carpal tunnel surgery. Ice was applied continually for 20-minute periods during the first three post-operative days. Hochberg found that patients using ice continually had a significantly greater decrease in pain. Unfortunately, the mode of ice application was not the same between the two groups, undermining the conclusions that can be drawn from the study. Bleakley *et al.*²⁴ compared two different regimes of ice treatment management in an attempt to assess the most effective. The use of standard ice application and the intermittent application of ice were employed amongst two different groups of patients presenting with acute ankle sprain. Their findings suggested that intermittent applications of ice after acute soft tissue injury enhanced the effects of ice in relation to pain relief on activity in the early stages (first week) of the injury, but made no significant difference in terms of function, swelling, or pain at rest. Their intermittent treatment regime involved 10 minutes of ice, 10 minutes off and 10 minutes of ice, repeated every two hours for the first 72 hours after injury, as recommended by MacAuley²⁵.

Coté *et al.*²⁶ investigated the use of ice, heat or a contrast bathing approach in first- and second-degree ankle sprains during the first three days after injury and before the onset of a rehabilitative exercise programme. This study concluded that the application of all three treatment modalities produced an increase in oedema in the post-acute phase in the sprained ankle; heat and contrast bathing produced identical increases in oedema; and ice therapy produced the least amount of oedema. The authors concluded that ice was the most appropriate treatment to use to minimise the development of oedema after injury.

The effects of barriers

The use of a barrier between the ice pack and skin is commonly recommended to protect the skin. Lavelle and Snyder²⁷ examined the effect of cooling when a barrier is present; they measured skin temperature after 30 minutes of ice application:

Type of barrier	Mean skin temperature
Padded bandage	30.5°C
Unpadded bandage	20.5°C
Dry washcloth	17.8°C
No barrier	10.8°C
Damp washcloth	9.9°C

The effect of subcutaneous fat acting as insulation has been examined by Hocutt²⁸ and Myrer²⁹. Hocutt *et al.* suggested that significant cooling occurred with ice application of 10 minutes to a depth of 2 centimetres (cm) in individuals with less than 1cm of fat. They suggested that in athletes with 2cm of fat, cooling was required for 20-30 minutes. Myrer and colleagues also showed that the depth of adipose tissue was a significant factor in the first 15 minutes of ice therapy, showing

an inverse relationship between adipose tissue and temperature decrease.

Ice and exercise

Bleakley *et al.*²⁴ in their systematic review concluded that there was marginal evidence for the combined use of ice and exercise.

Contraindications and adverse effects

Therapeutic cold treatment should be avoided on areas of reduced sensitivity in order to avoid the potential risk of frostbite. This can include, for example, patients with locally anaesthetised areas from nerve root compression syndromes, diabetic patients, patients with circulatory disorders, open wounds or skin conditions, patients with iron deficiency anaemia, and poor kidney function. Patients with a sensitivity to cold, such as those with Raynaud's syndrome and thyroid conditions, should also avoid cold treatment. Similarly, patients with severe cardiovascular disease, severe hypertension and those whose joint symptoms are aggravated by use of cold should all avoid cold application³⁰.

Frostbite is one of the most common, yet rare, adverse effects of cold application^{14,31}. Careless application of coolant sprays, particularly those containing ethyl chloride, are capable of causing localised frostbite^{32,33}.

Areas where nerves become more superficial should also be treated with extreme care and local cold application avoided; these include, for example, the ulnar nerve at the elbow and the peroneal nerve at the head of the fibular. Other cases in the literature have included cold-induced neuropathy in the axillary nerve and lateral femoral cutaneous neuropathy³⁴.

In common with so many other areas of healthcare, the evidence can be contradictory at times. The most recent work by Bleakley *et al.*²⁴ appears to show the most extensive consideration of the literature and, most helpfully, compares the various components of management for acute injuries.

References

1. Rivenburgh DW. Physical modalities in the treatment of injuries. *Clinical Sports Medicine*. 1992;11:645-959.
2. Meeusen R, Lievens P. The use of cryotherapy in sports injuries. *Sports Med*. 1986;3(6):398-414.
3. Swenson C, Swärd L, Karlsson J. Cryotherapy in sports medicine. *Scan J Med Sci Sports*. 1996;6(4):193-200.
4. Hochberg J. A randomised prospective study to assess the efficacy of two cold therapy treatments following carpal tunnel release. *J Hand Ther*. 2001;14:208-15.
5. Ksiezopolska-Pietrzak K. Cryotherapy in the treatment of rheumatic disease. *Ortop Traumatol Rehabil*. 2000;2(4):66-9.
6. d'Young AI. Domiciliary application of CryoCuff in severe haemophilia: qualitative questionnaire and clinical audit. *Haemophilia*. 2008;May 7 (Epub ahead of print).
7. Marek J, Jezdinsky J, Ochonsky P. Effects of local cold and heat therapy. *Acta Univ Palacki Olumuc Fac Med*. 1973;66:203-228.
8. Weston M, Taber C, Casagrande L *et al*. Changes in local blood volume during cold gel pack application to traumatised ankles. *Journal of Orthopaedics and Sports Physical Therapy (JOSPT)*. 1994;19(4):197-199.
9. Curl WW, Smith BP, Marr A *et al*. The effect of contusion and cryotherapy on

- skeletal muscle microcirculation. *Journal of Sports Medicine and Physical Fitness*. 1997;37(4):279-86.
10. Ho SS, Coel MN, Kagowa R et al. The effects of ice on blood flow and bone metabolism in knees. *Am J Sports Med*. 1994;22:537-540.
 11. Grana WA, Curl; WL, Reider B. Cold modalities. In: Drez D, ed. *Therapeutic modalities for Sports Injuries*. St Louis: Mosby Year Book; 1989:25-32.
 12. Knight KL. *Cryotherapy in Sport Injury Management*. Champaign, IL: Human Kinetics Publishers, 1995.
 13. Loeser JD. *Common Pain Problems: Guide to Practical Management*. Darien, Conn: Health Communications Inc; 1998:11-13.
 14. Harviksen K. Ice therapy in spasticity. *Acta Neurol Scand*. 1962;38:79-84.
 15. Lessard LA, Scudds RA, Amendola A et al. The efficacy of cryotherapy following arthroscopic knee surgery. *J Orthop Sports Phy Ther*. 1997;26:14-22.
 16. Hubbard TJ, Aronson SL, Denegar CR. Does cryotherapy hasten return to participation? *Journal of Athletic Training*. 2004;39(1):88-94.
 17. Pedowitz RN. Use of osteopathic manipulative treatment for iliotibial band friction syndrome. *J Am Osteopath Assoc*. 2005;105(12):563-7.
 18. Bleakley CM, McDonough SM, MacAuley DC. Cryotherapy for acute ankle sprains: a randomised controlled study of two different icing protocols. *British Journal of Sports Medicine*. 2006; 40:700-705.
 19. Chesterton LS, Foster NE, Ross L. Skin temperature response to cryotherapy. *Arch Phys Med Rehabil*. 2002;83:543-549.
 20. Zemke JE, Andersen JC, Guion WK et al. Intramuscular temperature response in the human leg to two forms of cryotherapy: ice massage and ice bag. *JOSPT*. 1998;27(4):301-307.
 21. Bilik SE (ed). *The Trainers Bible*. New York: TJ Reed and Co, 1946:257-263.
 22. Bevan R. *The Athletic Trainers Handbook*. Englewood Cliffs NJ: Prentice Hall, 1956:63-75.
 23. Stitik TP, Nadler SF. When – and how – to use cold most effectively. *Consultant*. 1998;38(12):2881-7.
 24. Bleakley C, McDonough S, MacAuley D. The use of ice in the management of acute soft-tissue injury: A systematic review of randomised controlled trials. *The American Journal of Sports Medicine*. 2004;32:251-261.
 25. MacAuley D. Ice therapy: how good is the evidence? *International Journal of Sports Medicine*. 2001;22:379-384.
 26. Côté DJ, Prentice WE, Hooker DN et al. Comparison of three ankle treatment procedures for minimising ankle sprain swelling. *Physical Therapy*. 1988;68(7):1072-1076.
 27. Lavelle BE, Snyder M. Differential conduction of cold through barriers. *J Adv Nur*. 1985;10:55-61.
 28. Hocutt JE, Jaffe R, Rylander CR et al. Cryotherapy in ankle sprains. *American Journal of Sports Medicine*. 1982;10:316-319.
 29. Myrer JW, Draper DO, Durrant E. Contrast therapy and intramuscular temperature in the human leg. *J Athl Train*. 1994;29:318-322.
 30. Williams A. *Spa Bodywork: A Guide for Massage Therapists*. Lippincott Williams & Wilkins, USA, 2006.
 31. Knight KL. Ice, compression, and elevation for initial injury care. In: Knight KL. *Cryotherapy: Theory, Technique and Physiology*. Chattanooga, Tenn: Chattanooga Corporation; 1985.
 32. Collins K, Storey M, Peterson K. Peroneal nerve palsy after cryotherapy.

Physician Sports Med. 1986;14:105-108.

33. Drez D, Faust DC, Evans JE. Cryotherapy and nerve palsy. *Am J Sports Med.* 1981;9:256-257.

34. Bassett FH, Kirkpatrick JS, Engelhardt DL *et al.* Cryotherapy induced nerve injury. *Am J Sports Med.* 1992;20:516-518.