



# Isothermic and fixed intensity heat acclimation methods induce similar heat adaptation following short and long-term timescales

Oliver R. Gibson<sup>a,\*</sup>, Jessica A. Mee<sup>a</sup>, James A. Tuttle<sup>b</sup>, Lee Taylor<sup>b</sup>, Peter W. Watt<sup>a</sup>, Neil S. Maxwell<sup>a</sup>

<sup>a</sup> Centre for Sport and Exercise Science and Medicine (SESAME), Exercise in Extreme Environments Laboratory, University of Brighton, Welkin Human Performance Laboratories, Denton Road, Eastbourne, UK

<sup>b</sup> Muscle Cellular and Molecular Physiology (MCMP) and Applied Sport and Exercise Science (ASEP) Research Groups, Department of Sport Science and Physical Activity, Institute of Sport and Physical Activity Research (ISPAR), University of Bedfordshire, Bedford Campus, Polhill Avenue, Bedfordshire, UK

## ARTICLE INFO

### Article history:

Received 17 December 2014

Received in revised form

11 February 2015

Accepted 11 February 2015

Available online 12 February 2015

### Keywords:

Heat illness

Heat stress

Hyperthermia

Taper

Temperature

Thermoregulation

## ABSTRACT

Heat acclimation requires the interaction between hot environments and exercise to elicit thermoregulatory adaptations. Optimal synergism between these parameters is unknown. Common practise involves utilising a fixed workload model where exercise prescription is controlled and core temperature is uncontrolled, or an isothermic model where core temperature is controlled and work rate is manipulated to control core temperature.

Following a baseline heat stress test; 24 males performed a between groups experimental design performing short term heat acclimation (STHA; five 90 min sessions) and long term heat acclimation (LTHA; STHA plus further five 90 min sessions) utilising either fixed intensity (50%  $\dot{V}_{O_{2peak}}$ ), continuous isothermic (target rectal temperature 38.5 °C for STHA and LTHA), or progressive isothermic heat acclimation (target rectal temperature 38.5 °C for STHA, and 39.0 °C for LTHA). Identical heat stress tests followed STHA and LTHA to determine the magnitude of adaptation.

All methods induced equal adaptation from baseline however isothermic methods induced adaptation and reduced exercise durations (STHA = −66% and LTHA = −72%) and mean session intensity (STHA = −13%  $\dot{V}_{O_{2peak}}$  and LTHA = −9%  $\dot{V}_{O_{2peak}}$ ) in comparison to fixed ( $p < 0.05$ ). STHA decreased exercising heart rate (−10 b min<sup>−1</sup>), core (−0.2 °C) and skin temperature (−0.51 °C), with sweat losses increasing (+0.36 L h<sup>−1</sup>) ( $p < 0.05$ ). No difference between heat acclimation methods, and no further benefit of LTHA was observed ( $p > 0.05$ ). Only thermal sensation improved from baseline to STHA (−0.2), and then between STHA and LTHA (−0.5) ( $p < 0.05$ ). Both the continuous and progressive isothermic methods elicited exercise duration, mean session intensity, and mean  $T_{rec}$  analogous to more efficient administration for maximising adaptation.

Short term isothermic methods are therefore optimal for individuals aiming to achieve heat adaptation most economically, i.e. when integrating heat acclimation into a pre-competition taper. Fixed methods may be optimal for military and occupational applications due to lower exercise intensity and simplified administration.

© 2015 Elsevier Ltd. All rights reserved.

## 1. Introduction

Repeated exposure to stressful hot environments initiates the heat-adapted phenotype. The heat-adapted phenotype is acquired

most effectively when hot and humid environmental conditions and physical work (intensity, duration and frequency) interact to stress thermoregulatory and cardiovascular systems (Sawka et al., 2011); this process is known as heat acclimation (Garrett et al., 2011). Primary adaptations induced by heat acclimation include decreased core temperature (Armstrong and Maresh, 1991; Buono et al., 1998; Garrett et al., 2011) and reduced heat storage (Aoyagi et al., 1997) facilitated by increased sudomotor function (Chinevere et al., 2008; Lorenzo and Minson, 2010; Machado-Moreira et al., 2006; Martinez et al., 2012), increased skin blood flow (Lorenzo and Minson, 2010), and cardiovascular adjustments eliciting greater maintenance of stroke volume and reduced heart rate at a

**Abbreviations:** FIXED, fixed intensity heat acclimation experimental group; HR, heart rate; HST, heat Stress Test; ISO<sub>CONT</sub>, continuous isothermic heat acclimation experimental group; ISO<sub>PROG</sub>, progressive isothermic heat acclimation experimental group; LTHA, long term heat acclimation; STHA, short term heat acclimation;  $T_{rec}$ , rectal temperature;  $T_{sk}$ , skin temperature;  $\dot{V}_{O_{2peak}}$ , peak oxygen uptake  
\* Corresponding author.

E-mail address: [o.r.gibson@brighton.ac.uk](mailto:o.r.gibson@brighton.ac.uk) (O.R. Gibson).

given workload (Frank et al., 2001). These adaptations contribute to a decreased thermal and perceptual strain (Castle et al., 2011), ultimately facilitating increased exercise performance in hot and cool environments (Lorenzo et al., 2010). Heat acclimation is often categorised into short term (STHA) and long term (LTHA) induction periods. LTHA, the traditional time scale, generally comprises  $\geq 10$  daily heat exposures (Garrett et al., 2011), potentiating the most complete phenotypic adaptation. STHA utilises  $\leq 5$  daily exposures, facilitating rapid, but, incomplete adaptation ( $\sim 75\%$  compared to LTHA (Pandolf, 1979)). Notwithstanding, STHA still remains an effective tool used by practitioners for augmenting adaptation before exposure to hot environments, improving tolerance to exercise or work (Garrett et al., 2012, 2009).

Increased core temperature is a fundamental requirement for inducing heat acclimation (Regan et al., 1996; Taylor and Cotter, 2006). Isothermic heat acclimation (also known as controlled hyperthermia) is imposed based upon endogenous (internal) criteria (Castle et al., 2012; Garrett et al., 2014, 2012, 2009; Hom et al., 2012; Machado-Moreira et al., 2006; Magalhães et al., 2010a, 2010b; Patterson et al., 2014, 2004), and might provide sustained targeting and attainment of specific and individualised internal temperatures through a combination of active and passive heat acclimation (Fox et al., 1963). The balance between work and rest to target and maintain specific core temperatures ensures a consistency, or a progression of endogenous heat strain to induce adaptation, albeit requiring alterations in administration throughout each session. Implementation of fixed intensity heat acclimation methods is in comparison relatively simple, with participants maintaining a fixed workload throughout each active acclimation session (Amorim et al., 2011; Castle et al., 2011; Cheung and McLellan, 1998; Houmard et al., 1990; Kresfelder et al., 2006; Lorenzo and Minson, 2010; Lorenzo et al., 2010; Marshall et al., 2007; Nielsen et al., 1997, 1993; Sandström et al., 2008; Watkins et al., 2008; Yamada et al., 2007). Fixed methods derive exercise workloads from a pre acclimation baseline, and the exogenous (external) environment are consistent day-on-day. Though this method may provide sufficient heat strain during the initial sessions of heat acclimation regimens, fixed methods may not achieve the desired, nor optimally potentiating stimuli – increased core temperature, as the thermal strain relative to the start of acclimation diminishes with ensuing adaptation (Taylor and Cotter, 2006; Taylor, 2014). During both STHA and LTHA, relative workload and the thermal strain of heat acclimation are likely to reduce during fixed intensity as on-going adaptation is seen. Isothermic heat acclimation, where endogenous thermal stimulus is consistently targeted throughout, may positively sustain the rate of adaptation, or advance adaptation should a progressive increase in core temperature be implemented (Taylor and Cotter, 2006; Taylor, 2014). Progressive isothermic methods have only previously been implemented using models where the environmental conditions or workload for acclimation are increased (Burk et al., 2012; Chen et al., 2013; Daanen et al., 2011), this presumably to offset the aforementioned ongoing adaptation. These progressive methods are not certain to increase core temperature in the manner that a progressive increase in the isothermic target temperature would. Varied administration of heat acclimation methods has likely produced different phenotypic adaptive responses. The mode of exercise, relative exercise intensity and climatic conditions may modulate different degrees of adaptation (Taylor and Cotter, 2006). Should the anticipated core temperature changes be observed between methods it is likely that fixed heat acclimation methods are analogous to a reduction in the potentiating stimuli for adaptation and consequently the rate of adaptation would decrease from STHA to LTHA. The isothermic continuous method should theoretically sustain potentiating stimuli and consequently sustain the rate of adaptation from STHA to LTHA.

Finally a progressive isothermic method could theoretically be used to increase potentiating stimuli and may increase the rate of adaptation from STHA to LTHA.

The aim of the present study was to determine whether any differences in heat adaptation occurred between an established exogenous controlled, fixed intensity heat acclimation method, an endogenous controlled, isothermic heat acclimation method, and a stepwise progressive endogenous isothermic heat acclimation method, after STHA and LTHA periods. No direct comparison has been made of the observed adaptation and administration differences between isothermic and fixed heat acclimation methods across STHA and LTHA timescales; additionally evidence is limited in support of a stepwise progression in thermal strain to increase the rate of adaptation from STHA to LTHA. We hypothesised that the rate of phenotypic adaptation would be greater in isothermic heat acclimation methods in comparison to fixed methods due to sustained strain. It was additionally hypothesised that a greater rate of adaptation would be induced by utilising a progressive model. It was also hypothesised that implementation of isothermic heat acclimation would require reduced exercise durations and lower average sessional exercise intensities, in spite of initially higher exercise intensities, which would favour athletes in the pre-competition taper.

## 2. Methods

### 2.1. Participants

Twenty-four healthy males were assigned into fixed intensity (FIXED), or isothermic heat acclimation (ISO) groups, ISO was then subdivided into continuous isothermic heat acclimation (ISO<sub>CONT</sub>), or progressive isothermic heat acclimation (ISO<sub>PROG</sub>) groups; participants were matched for peak oxygen uptake ( $\dot{V}_{O_{2peak}}$ ) and anthropometric characteristics. Data are presented in Table 1. Confounding variables of smoking, caffeine, glutamine, alcohol, generic supplementation, prior thermal, hypoxic and hyperbaric exposures were all controlled in line with previous work in the field (Gibson et al., 2014; Taylor et al., 2011). Following institutional ethics approval and full description of experimental procedures, all participants completed medical questionnaires and provided written informed consent following the principles outlined by the declaration of Helsinki of 1975, as revised in 2013. The experimental design for the study is presented in Fig. 1 with full explanation of the heat acclimation methods contained within the “Heat Acclimation Methods” Section 2.4 which follows.

### 2.2. Preliminary testing

Participants consumed 500 mL of water 2 h before all preliminary and experimental exercise sessions (Sawka et al., 2007). A urine osmometer (Alago Vitech Scientific, Pocket PAL-OSMO, UK) was used to ensure consistent hydration prior to each experimental session (Garrett et al., 2014). Participants were deemed euhydrated and subsequently able to commence further preliminary, and experimental procedures if urine osmolality was  $< 700 \text{ mOsm kg}^{-1} \text{ H}_2\text{O}$  (Sawka et al., 2007). Prior to the initial  $\dot{V}_{O_{2peak}}$  experimental trial, height (cm) using a fixed stadiometer (Detecto Physicians Scales; Cranlea & Co., Birmingham, UK), and body density, using calipers (Harpender, Burgess Hill, UK) and a four site skin fold calculation (Durnin and Womersley, 1974) were determined, later body fat (%) was calculated from body density (Siri, 1956) and body surface area (Du Bois and Du Bois, 1916). Nude body mass (NBM) was recorded to 0.01 kg from digital scales (ADAM GFK 150, USA), relative metabolic heat production (MHP;

**Table 1**

Mean  $\pm$  SD Participant characteristics and descriptive data for the initial workload in fixed intensity heat acclimation (FIXED), continuous isothermic heat acclimation (ISO<sub>CONT</sub>), and progressive isothermic heat acclimation (ISO<sub>PROG</sub>) experimental groups.

	FIXED (n=8)	ISO <sub>CONT</sub> (n=8)	ISO <sub>PROG</sub> (n=8)
Age (years)	19.9 $\pm$ 1.0	22.6 $\pm$ 5.5	26.1 $\pm$ 4.9*
Height (cm)	179.3 $\pm$ 5.8	177.9 $\pm$ 5.8	179.5 $\pm$ 6.6
Body mass (kg)	79.2 $\pm$ 18.3	74.2 $\pm$ 6.9	75.1 $\pm$ 8.8
Body mass index (kg m <sup>2</sup> )	24.6 $\pm$ 5.7	23.4 $\pm$ 1.7	23.4 $\pm$ 3.2
Body surface area (m <sup>2</sup> )	1.97 $\pm$ 0.21	1.92 $\pm$ 0.11	1.94 $\pm$ 0.11
Body fat (%)	14.9 $\pm$ 7.7	14.8 $\pm$ 2.2	14.1 $\pm$ 3.5
$\dot{V}O_{2peak}$ (L min <sup>-1</sup> )	3.61 $\pm$ 0.90	3.63 $\pm$ 0.69	3.80 $\pm$ 0.55
Initial workload (W kg <sup>-1</sup> )	1.6 $\pm$ 0.5	2.2 $\pm$ 0.3*	2.4 $\pm$ 0.4*
Initial metabolic heat production (W kg <sup>-1</sup> )	9.5 $\pm$ 3.3	11.1 $\pm$ 1.4	11.9 $\pm$ 2.0

\* Significantly difference from FIXED ( $p < 0.05$ ).

(W kg<sup>-1</sup>) was calculated to describe the initial requirements of each heat acclimation method in accordance with the guidelines of Cramer and Jay (2014).

$\dot{V}O_{2peak}$  (L min<sup>-1</sup>) was determined from an incremental test on a cycle ergometer (Monark e724, Vansbro, Sweden) in temperate laboratory conditions (20 °C, 40% relative humidity (RH)). Starting intensity was set at 80 W, with resistance applied to the flywheel eliciting a 24 W min<sup>-1</sup> increase at the constant cadence of 80 rpm. Expired metabolic gas was measured using online gas analysis (Metamax 3X, Cortex, Germany);  $\dot{V}O_{2peak}$  was considered as the highest  $\dot{V}O_2$  obtained in any 10 s period. Heart rate (HR; b min<sup>-1</sup>) was recorded continually during all exercise tests by telemetry (Polar Electro Oyo, Temple, Finland). Saddle position was adjusted by the participant to their preferred cycling position and remained unchanged for all experimental trials. Heat acclimation workloads were subsequently calculated using linear regression utilising power:  $\dot{V}O_2$  data collected following the incremental test.

### 2.3. Heat stress testing


A running heat stress test (HST) was performed as a preliminary test (HST1), then 48 h after STHA (HST2), and 48 h after LTHA (HST3) inside a purpose built environmental chamber with temperature and humidity (40.0  $\pm$  0.1 °C and 28.4  $\pm$  6.6% RH)

controlled using automated computer feedback (WatFlow control system; TISS, Hampshire, UK) and no additional convective cooling provided e.g. motorised fan. All HST were performed between 08:00 and 10:00 h (Drust et al., 2005). Following confirmation of adequate hydration, participants inserted a single-use disposable rectal thermistor (Henleys Medical, UK, Meter logger Model 401, Yellow Springs Instruments, Yellow Springs, Missouri, USA; accuracy  $\pm$  0.20 °C) 10 cm past the anal sphincter to measure rectal temperature ( $T_{rec}$ ). Skin temperature ( $T_{sk}$ ) was measured using a data logger (Squirrel Meter Logger, Grant Instruments, Cambridge, UK) and skin thermistors attached to the right-hand side of the body using zinc oxide tape (Cramer Products Inc., Kansas, USA) at the pectoralis major muscle belly, lateral head of triceps brachii, rectus femoris muscle belly and lateral head of the gastrocnemius (Ramanathan, 1964). Mean skin temperature was calculated according to the formula of Ramanathan (1964). Absolute sweat loss (L h<sup>-1</sup>) was estimated using the change in towel-dried NBM from the pre-to-post exercise periods and adjusted based upon the HST duration. Participants were not permitted to consume any fluid between pre and post-test measurement of NBM. No correction was made for insensible water loss and loss of mass associated with the respiratory exchange of O<sub>2</sub> and CO<sub>2</sub> (Dion et al., 2013); all were assumed to be similar between HSTs due to the equal length of each trial at each time point (Table 2).

After a 20 min seated stabilisation period in temperate laboratory conditions, resting measures were taken after which participants entered the environmental chamber to perform 30 min running at 9 km h<sup>-1</sup> and 2% elevation. HR,  $T_{rec}$  and  $T_{sk}$  were recorded every 5 min. Ratings of perceived exertion (Borg et al., 1985) and thermal sensation (Toner et al., 1986) were recorded every 10 min. HSTs were terminated if  $T_{rec} \geq 39.7$  °C (zero incidences), or the participant withdrew due to volitional exhaustion, or inability to maintain the running speed despite strong verbal encouragement.

### 2.4. Heat acclimation methods

Each heat acclimation testing session was conducted at the same time of day (07:00–11:00 h) to control for effects of daily variation in performance (Shido et al., 1999). Following provision of a urine sample and NBM, each participant inserted the rectal thermistor described in the HST and affixed a HR monitor upon which time resting measures were taken after 5 min seated in temperate laboratory conditions. Participants subsequently



**Fig. 1.** Experimental schematic. See text for details.

**Table 2**

Mean  $\pm$  SD Protocol, thermoregulatory and physiological response data for STHA, then LTHA of fixed intensity heat acclimation (FIXED), continuous isothermic heat acclimation (ISO<sub>CONT</sub>), and progressive isothermic heat acclimation (ISO<sub>PROG</sub>) experimental groups.

	STHA			LTHA		
	FIXED	ISO <sub>CONT</sub>	ISO <sub>PROG</sub>	FIXED	ISO <sub>CONT</sub>	ISO <sub>PROG</sub>
Exercising duration (min)	450 $\pm$ 0	337 $\pm$ 47 <sup>#</sup>	263 $\pm$ 47 <sup>†</sup>	900 $\pm$ 0 <sup>*</sup>	707 $\pm$ 102 <sup>* #</sup>	598 $\pm$ 87 <sup>* †</sup>
Time to target $T_{rec}$ (min)	89 $\pm$ 2	49 $\pm$ 12 <sup>#</sup>	43 $\pm$ 10 <sup>#</sup>	89 $\pm$ 2	54 $\pm$ 14 <sup>* #</sup>	52 $\pm$ 10 <sup>* #</sup>
Mean session intensity (% $\dot{V}_{O_{2peak}}$ )	50.0 $\pm$ 0.0	40.6 $\pm$ 6.2 <sup>#</sup>	33.5 $\pm$ 7.2 <sup>#</sup>	50.0 $\pm$ 0.0	43.9 $\pm$ 6.0 <sup>*</sup>	38.1 $\pm$ 7.1 <sup>* #</sup>
Mean exercise intensity (% $\dot{V}_{O_{2peak}}$ )	50.0 $\pm$ 0.0	55.2 $\pm$ 6.4	57.7 $\pm$ 5.3	50.0 $\pm$ 0.0	56.7 $\pm$ 5.8	57.5 $\pm$ 4.4
Mean session power (W kg <sup>-1</sup> )	1.6 $\pm$ 0.4	1.4 $\pm$ 0.1	1.3 $\pm$ 0.3	1.6 $\pm$ 0.4	1.5 $\pm$ 0.2 <sup>*</sup>	1.4 $\pm$ 0.3 <sup>*</sup>
Mean exercise power (W kg <sup>-1</sup> )	1.6 $\pm$ 0.4	1.9 $\pm$ 0.3	2.2 $\pm$ 0.5 <sup>#</sup>	1.6 $\pm$ 0.4	2.0 $\pm$ 0.2	2.2 $\pm$ 0.5 <sup>#</sup>
Total work done (kJ)	3352 $\pm$ 815	2789 $\pm$ 358	2590 $\pm$ 560	6701 $\pm$ 1603 <sup>*</sup>	6113 $\pm$ 834 <sup>*</sup>	5880 $\pm$ 1484 <sup>*</sup>
Mean $T_{rec}$ (°C)	38.03 $\pm$ 0.16	38.18 $\pm$ 0.12	38.26 $\pm$ 0.18 <sup>*</sup>	37.92 $\pm$ 0.15 <sup>*</sup>	38.16 $\pm$ 0.12 <sup>#</sup>	38.26 $\pm$ 0.20 <sup>#</sup>
Mean $T_{recfinal60}$ (°C)	38.28 $\pm$ 0.18	38.44 $\pm$ 0.13	38.60 $\pm$ 0.18 <sup>#</sup>	38.16 $\pm$ 0.18 <sup>*</sup>	38.43 $\pm$ 0.14 <sup>#</sup>	38.63 $\pm$ 0.21 <sup>#</sup>
$\Delta T_{rec}$ (°C)	1.72 $\pm$ 0.58	1.74 $\pm$ 0.19	2.15 $\pm$ 0.38	1.63 $\pm$ 0.60	1.78 $\pm$ 0.22	2.27 $\pm$ 0.29 <sup>#</sup>
Rate $T_{rec}$ increase (°C h <sup>-1</sup> )	1.43 $\pm$ 0.43	1.98 $\pm$ 0.31	2.42 $\pm$ 0.54	1.34 $\pm$ 0.46	1.99 $\pm$ 0.36	2.36 $\pm$ 0.50
Duration $T_{rec} \geq 38.5$ °C (min)	118 $\pm$ 53	161 $\pm$ 62	244 $\pm$ 62 <sup>†</sup>	176 $\pm$ 86 <sup>*</sup>	318 $\pm$ 118 <sup>* #</sup>	462 $\pm$ 120 <sup>* †</sup>
Duration $T_{rec} \geq 39.0$ °C (min)	17 $\pm$ 21	5 $\pm$ 14 <sup>+</sup>	38 $\pm$ 32	21 $\pm$ 24	13 $\pm$ 24	146 $\pm$ 70 <sup>* †</sup>
Mean HR (b min <sup>-1</sup> )	155 $\pm$ 13	150 $\pm$ 9	142 $\pm$ 11	150 $\pm$ 13	150 $\pm$ 9	143 $\pm$ 11

\* Significantly different ( $p < 0.05$ ) from STHA (within group).

# Significantly different ( $p < 0.05$ ) from FIXED (within timescale).

† Significantly different ( $p < 0.05$ ) from FIXED and ISO<sub>CONT</sub> (within timescale).

+ Significantly different ( $p < 0.05$ ) from ISO<sub>PROG</sub> within timescale.

mounted a cycle ergometer (Monark, e724, Vansbro, Sweden) located inside the environmental chamber where conditions were consistent for all groups ( $40.2 \pm 0.4$  °C,  $39.0 \pm 7.8\%$  RH). The FIXED participants performed 10 sessions of 90 min of continuous cycling exercise at a workload corresponding to 50%  $\dot{V}_{O_{2peak}}$ . ISO<sub>CONT</sub> participants exercised initially at a workload corresponding to 65% of  $\dot{V}_{O_{2peak}}$  until a target  $T_{rec}$  of 38.5 °C was achieved for all 10 heat acclimation sessions. ISO<sub>PROG</sub> participants exercised initially at a workload corresponding to 65% of  $\dot{V}_{O_{2peak}}$  targeting a  $T_{rec}$  of 38.5 °C for the first five sessions, then progressing to a  $T_{rec}$  of 39.0 °C for the final five sessions. Once target  $T_{rec}$  had been reached, power was adjusted every 5 min, first by a 25%  $\dot{V}_{O_{2peak}}$  reduction and then adjusted ( $\pm 5\%$   $\dot{V}_{O_{2peak}}$ , or seated rest) to maintain the desired experimental  $T_{rec}$  for a total session duration of 90 min within the environmental chamber; workloads at the onset of exercise are presented in Table 1. During each testing session HR,  $T_{rec}$  and power output, were recorded every 5 min. Mean  $T_{rec}$  reflects the average  $T_{rec}$  recorded throughout each acclimation method.  $T_{recfinal60-min}$  quantifies the mean  $T_{rec}$  between minutes 30 and 90 of the acclimation session to reflect the temperature following the initial rate of increase. Exercising duration was defined as the total time exercising (power output  $> 1$  W) during acclimation sessions reflecting the physical work demands throughout each 90 min session. Mean Session Intensity (%  $\dot{V}_{O_{2peak}}$  and W kg<sup>-1</sup>) was calculated from the relative exercise intensity during each 5 min period throughout all of the 90 min acclimation sessions. This contrasted the Mean Exercise Intensity (%  $\dot{V}_{O_{2peak}}$  and W kg<sup>-1</sup>), which reflected the mean relative exercise intensity only (power output  $> 1$  W), thus excluding periods of rest within isothermic methods. Data for these variables are provided in Table 2.

## 2.5. Statistical analyses

All outcome variables were first checked for normality using Kolmogorov–Smirnov and sphericity using the Greenhouse Geisser method prior to further analysis. Two way mixed design ANOVA were performed to determine differences in dependent variables between heat acclimation methods for STHA and LTHA timescales, and between heat acclimation methods and HST1,

HST2 and HST3. Adjusted Bonferroni comparisons were used as post hoc analyses, determining where differences existed within ANOVA where a time or interaction was found. Data are reported as mean  $\pm$  SD, with two-tailed significance was accepted at  $p < 0.05$ .

## 3. Results

### 3.1. Method administration and thermoregulatory and physiological responses to short and long term heat acclimation

Differences ( $p < 0.05$ ) were observed with increased exercise duration, total work done and duration  $T_{rec} \geq 38.5$  °C in all methods from STHA to LTHA, data are presented in Table 2. Mean session intensity, mean session power and the time to target  $T_{rec}$  increased from STHA to LTHA in ISO<sub>CONT</sub> and ISO<sub>PROG</sub>, with mean  $T_{rec}$  and mean  $T_{recfinal60}$ , reducing in FIXED. The duration  $T_{rec} \geq 39.0$  °C increased from STHA to LTHA in ISO<sub>PROG</sub> only. No difference was observed for mean exercise intensity ( $f=1.935$ ,  $p=0.179$ ), mean exercise power ( $f=1.061$ ,  $p=0.315$ ), change in  $T_{rec}$  ( $f=0.866$ ,  $p=0.363$ ), rate of  $T_{rec}$  increase ( $f=2.158$ ,  $p=0.157$ ), or mean HR ( $f=3.026$ ,  $p=0.097$ ) between STHA and LTHA.

A between heat acclimation methods interaction effect was observed for exercise duration ( $f=13.090$ ,  $p < 0.001$ ), and the time to target  $T_{rec}$  ( $f=6.500$ ,  $p=0.006$ ), mean session intensity ( $f=6.727$ ,  $p=0.006$ ), mean  $T_{rec}$  ( $f=7.063$ ,  $p=0.005$ ), mean  $T_{recfinal60}$  ( $f=11.073$ ,  $p=0.001$ ), duration  $T_{rec} \geq 38.5$  °C ( $f=14.608$ ,  $p < 0.001$ ), duration  $T_{rec} \geq 39.0$  °C ( $f=28.262$ ,  $p < 0.001$ ), mean exercise power ( $f=3.765$ ,  $p=0.040$ ), change in  $T_{rec}$  ( $f=5.277$ ,  $p=0.014$ ) and mean HR ( $f=11.073$ ,  $p=0.001$ ). Post-hoc analysis is presented in Table 2 for clarity. No between group interaction was observed for total work done ( $f=0.011$ ,  $p=0.989$ ), mean exercise intensity ( $f=3.186$ ,  $p=0.062$ ), rate of  $T_{rec}$  increase ( $f=0.884$ ,  $p=0.428$ ), or for the post-hoc analysis of mean session power ( $f=4.822$ ,  $p=0.019$ ).

### 3.2. Daily responses to heat acclimation

Resting  $T_{rec}$  ( $f=3.048$ ,  $p=0.002$ ), resting HR ( $f=3.085$ ,  $p=0.002$ ), and sessional sweat loss (% NBM) ( $f=3.798$ ,  $p < 0.000$ )



all demonstrated improvements overall as the number of heat acclimation sessions increased. Post-hoc analysis revealed resting  $T_{\text{rec}}$  was reduced ( $p < 0.05$ ) from session one before session eight, nine and 10, resting HR was reduced ( $p < 0.05$ ) from session one before session nine and 10 and sweat loss was increased ( $p < 0.05$ ) from session one following session eight, nine and 10. No between group effect was observed for resting  $T_{\text{rec}}$  ( $f=1.146$ ,  $p=0.311$ ), resting HR ( $f=1.553$ ,  $p=0.076$ ) and sessional sweat loss ( $f=1.007$ ,  $p=0.453$ ). Data are presented in Fig. 3.

### 3.3. Heat stress testing – resting adaptations

Resting HR ( $f=7.730$ ,  $p=0.001$ ) and resting  $T_{\text{rec}}$  ( $f=7.372$ ,  $p=0.004$ ) reduced with heat acclimation; post-hoc analysis revealed a reduction ( $p < 0.05$ ) in both measures from HST1 to HST2 and HST1 to HST3, but no difference between HST2 and HST3 ( $p > 0.05$ ), data are presented in Table 3. No between heat acclimation method interaction was observed for either resting HR ( $f=0.819$ ,  $p=0.521$ ) or resting  $T_{\text{rec}}$  ( $f=0.750$ ,  $p=0.537$ ).

### 3.4. Heat stress testing – exercising adaptations

Mean exercising HR ( $f=23.887$ ,  $p < 0.001$ ), mean  $T_{\text{rec}}$  ( $f=11.067$ ,  $p < 0.001$ ), sweat loss ( $f=10.516$ ,  $p < 0.001$ ), mean  $T_{\text{sk}}$  ( $f=10.516$ ,  $p < 0.001$ ) and peak  $T_{\text{sk}}$  ( $f=13.185$ ,  $p < 0.001$ ) reduced with heat acclimation; post-hoc analysis revealed a reduction ( $p < 0.05$ ) in mean exercising HR, mean  $T_{\text{rec}}$ , mean  $T_{\text{sk}}$ , and peak  $T_{\text{sk}}$ , and increase in sweat loss from HST1 to HST2 and HST1 to HST3, but no difference between HST2 and HST3 ( $p > 0.05$ ) (Table 3). No between heat acclimation method interaction was observed for mean exercising HR ( $f=0.431$ ,  $p=0.786$ ), mean  $T_{\text{rec}}$  ( $f=0.213$ ,  $p=0.930$ ), sweat loss ( $f=2.183$ ,  $p=0.870$ ) or peak  $T_{\text{sk}}$  ( $f=2.008$ ,  $p=0.111$ ). No changes were observed between HSTs, or between heat acclimation methods for exercise duration ( $f=2.333$ ,  $p=0.125$ ) and ( $f=0.333$ ,  $p=0.854$ ), change in exercising HR ( $f=0.529$ ,  $p=0.593$ ) and ( $f=2.318$ ,  $p=0.073$ ), the change  $T_{\text{rec}}$  ( $f=0.126$ ,  $p=0.295$ ) and ( $f=0.975$ ,  $p=0.432$ ), or the rate of  $T_{\text{rec}}$  increase ( $f=1.257$ ,  $p=0.295$ ) and ( $f=0.975$ ,  $p=0.432$ ) respectively.

### 3.5. Heat stress testing – perceptual changes

Peak thermal sensation ( $f=8.316$ ,  $p=0.001$ ) and mean thermal sensation ( $f=5.573$ ,  $p=0.007$ ) reduced with heat acclimation,

post-hoc analysis revealed a reduction ( $p < 0.05$ ) in peak thermal sensation from HST1 to HST2, HST1 to HST3, and HST2 to HST3 and mean thermal sensation from HST2 to HST3 (Table 3). No between heat acclimation method interaction was observed for either peak thermal sensation ( $f=1.137$ ,  $p=0.352$ ) or mean thermal sensation ( $f=1.150$ ,  $p=0.346$ ). No changes were observed between HSTs, or between heat acclimation methods for peak rating of perceived exertion ( $f=2.891$ ,  $p=0.067$ ) and ( $f=2.194$ ,  $p=0.086$ ), or mean rating of perceived exertion ( $f=1.787$ ,  $p=0.180$ ) and ( $f=0.705$ ,  $p=0.593$ ), respectively.

## 4. Discussion

The aim of this experiment was to determine whether there was a difference in measures of heat adaptation to STHA and LTHA between a fixed intensity heat acclimation method, a controlled isothermic heat acclimation method and a progressive isothermic heat acclimation method. It was observed that equal heat adaptation was induced between all methods over STHA with no significant additional benefit from our LTHA timescale. Relative to fixed intensity methods, Isothermic methods are the favourable form of administration with equal adaptation induced following reduced exercise durations and mean session intensity.

### 4.1. Differences in heat acclimation methods

No difference in the magnitude of adaptation existed between fixed intensity, continuous isothermic, and progressive isothermic heat acclimation methods, however during STHA and LTHA, the FIXED mode of heat acclimation was inferior to isothermic heat acclimation methods (ISO<sub>CONT</sub> and ISO<sub>PROG</sub>) when considering applied practical perspectives in accordance with established recommendations for interacting physical training and heat acclimation to maintain performance (Aoyagi et al., 1997). Isothermic methods achieved adaptation with reduced exercise durations (STHA and LTHA) and mean session intensity (STHA and LTHA), which is desirable for athletic applications as an effective means for reducing the volume of physiological strain of exercise in the heat. This application perhaps contrasts military and occupational applications for which the FIXED method may be optimal due to lower intensity of work and simplified administration facilitating implementation for large cohorts, or a research perspective when

**Table 3**

Mean  $\pm$  SD heat stress test data at baseline (HST1), post five HA sessions (HST2) and post 10 HA sessions (HST3) of the fixed intensity heat acclimation (FIXED), continuous isothermic heat acclimation (ISO<sub>CONT</sub>), and progressive isothermic heat acclimation (ISO<sub>PROG</sub>) experimental groups.

	HST1			HST2			HST3		
	FIXED	ISO <sub>CONT</sub>	ISO <sub>PROG</sub>	FIXED	ISO <sub>CONT</sub>	ISO <sub>PROG</sub>	FIXED	ISO <sub>CONT</sub>	ISO <sub>PROG</sub>
Duration (min)	25.0 $\pm$ 8.0	29.4 $\pm$ 1.8	29.4 $\pm$ 1.8	25.0 $\pm$ 8.0	30.0 $\pm$ 0.0	30.0 $\pm$ 0.0	25.6 $\pm$ 8.2	30.0 $\pm$ 0.0	30.0 $\pm$ 0.0
Rest HR (b min <sup>-1</sup> )	74 $\pm$ 8	71 $\pm$ 9	63 $\pm$ 10	65 $\pm$ 11*	66 $\pm$ 8*	59 $\pm$ 9*	69 $\pm$ 9*	63 $\pm$ 4*	56 $\pm$ 12*
Change HR (b min <sup>-1</sup> )	107 $\pm$ 9	113 $\pm$ 13	118 $\pm$ 11	115 $\pm$ 16	105 $\pm$ 12	111 $\pm$ 9	111 $\pm$ 11	110 $\pm$ 10	115 $\pm$ 10
Mean exercising HR (b min <sup>-1</sup> )	161 $\pm$ 10	159 $\pm$ 9	154 $\pm$ 17	152 $\pm$ 12*	147 $\pm$ 10*	145 $\pm$ 13*	153 $\pm$ 9*	148 $\pm$ 8*	144 $\pm$ 18*
Rest $T_{\text{rec}}$ (°C)	37.23 $\pm$ 0.35	37.05 $\pm$ 0.21	36.94 $\pm$ 0.40	36.94 $\pm$ 0.36*	36.95 $\pm$ 0.21*	36.73 $\pm$ 0.41*	36.90 $\pm$ 0.40*	36.96 $\pm$ 0.19*	36.75 $\pm$ 0.25*
Change $T_{\text{rec}}$ (°C)	1.18 $\pm$ 0.44	1.61 $\pm$ 0.31	1.48 $\pm$ 0.19	1.25 $\pm$ 0.57	1.39 $\pm$ 0.35	1.44 $\pm$ 0.25	1.20 $\pm$ 0.47	1.28 $\pm$ 0.38	1.41 $\pm$ 0.39
Rate $T_{\text{rec}}$ (°C h <sup>-1</sup> )	2.35 $\pm$ 0.87	3.21 $\pm$ 0.62	2.97 $\pm$ 0.39	2.49 $\pm$ 1.13	2.77 $\pm$ 0.71	2.87 $\pm$ 0.49	2.39 $\pm$ 0.94	2.56 $\pm$ 0.75	2.82 $\pm$ 0.78
Mean $T_{\text{rec}}$ (°C)	37.77 $\pm$ 0.30	37.76 $\pm$ 0.17	37.56 $\pm$ 0.39	37.58 $\pm$ 0.34*	37.57 $\pm$ 0.19*	37.41 $\pm$ 0.41*	37.45 $\pm$ 0.30*	37.52 $\pm$ 0.27*	37.35 $\pm$ 0.43*
Sweat loss (L h <sup>-1</sup> )	1.45 $\pm$ 0.50	1.61 $\pm$ 0.43	1.28 $\pm$ 0.42	1.88 $\pm$ 0.75*	1.95 $\pm$ 0.36*	1.96 $\pm$ 0.80*	2.16 $\pm$ 0.61*	2.17 $\pm$ 0.61*	1.73 $\pm$ 0.58*
Peak $T_{\text{sk}}$ (°C)	37.72 $\pm$ 0.98	37.40 $\pm$ 0.78	37.52 $\pm$ 0.48	36.95 $\pm$ 1.11*	36.74 $\pm$ 0.32*	36.81 $\pm$ 0.84*	37.16 $\pm$ 0.83*	36.73 $\pm$ 0.62*	36.73 $\pm$ 0.81*
Mean $T_{\text{sk}}$ (°C)	35.70 $\pm$ 1.07	36.07 $\pm$ 0.67	36.05 $\pm$ 0.52	35.47 $\pm$ 1.12*	35.65 $\pm$ 0.38*	35.38 $\pm$ 0.86*	35.74 $\pm$ 0.99*	34.93 $\pm$ 0.71*	35.13 $\pm$ 0.77*
Peak RPE	13 $\pm$ 3	16 $\pm$ 3	15 $\pm$ 2	14 $\pm$ 4	15 $\pm$ 3	14 $\pm$ 3	14 $\pm$ 3	14 $\pm$ 3	14 $\pm$ 4
Mean RPE	13 $\pm$ 3	14 $\pm$ 3	13 $\pm$ 1	13 $\pm$ 3	14 $\pm$ 3	13 $\pm$ 2	13 $\pm$ 3	13 $\pm$ 3	13 $\pm$ 2
Peak TSS	7.1 $\pm$ 1.0	6.9 $\pm$ 0.8	6.7 $\pm$ 0.5	6.8 $\pm$ 0.8	6.7 $\pm$ 1.0	6.7 $\pm$ 0.4	6.5 $\pm$ 1.0*	6.3 $\pm$ 1.0*	6.3 $\pm$ 0.5*
Mean TSS	6.8 $\pm$ 1.0	6.3 $\pm$ 0.6	6.3 $\pm$ 0.4	6.5 $\pm$ 0.9	6.3 $\pm$ 0.8	6.3 $\pm$ 0.5	6.0 $\pm$ 0.8*	6.1 $\pm$ 0.5*	6.2 $\pm$ 1.0*

\* HST significantly different ( $p < 0.05$ ) from HST1 overall.

# HST significantly different ( $p < 0.05$ ) from HST2 overall.

matching of training stimulus is required. Exercising durations were lower in isothermic methods (66–79% of session duration) compared to fixed intensity heat acclimation (>99% of session duration). It is noteworthy that the time taken to achieve the target  $T_{\text{rec}}$  in ISO<sub>CONT</sub> and ISO<sub>PROG</sub> increased from STHA to LTHA by 6.6% and 10.0% respectively, thus demonstrating the effects of ongoing adaptation, and using the ISO<sub>PROG</sub> method the greater work required to attain a higher  $T_{\text{rec}}$ . Higher initial work intensity balanced by increased rest periods are congruous with typical training regimes, therefore isothermic methods may be more appropriate when integrating heat acclimation into a training taper (Mujika et al., 2004) prior to competition due to sport specificity (Houmard et al., 1990), particularly when acknowledging that the duration of a typical fixed heat acclimation session is at the upper end of that desirable for typical pre-competition training session, with the repeated sessions exceeding the typical volume of a typical endurance training taper (Spilsbury et al., 2014).

#### 4.2. Adaptations made during short and long term heat acclimation


Isothermic heat acclimation methods were more favourable than FIXED at targeting and sustaining specific  $T_{\text{rec}}$  (i.e.  $\geq 38.5^\circ\text{C}$ ) thus delivering greater elevations in thermal strain, notably the important potentiating stimuli of increased core temperature over both STHA and LTHA (Regan et al., 1996; Taylor and Cotter, 2006). This statement can be evidenced by ISO<sub>CONT</sub> and ISO<sub>PROG</sub> evoking greater mean  $T_{\text{rec}}$ , mean  $T_{\text{rec final 60}}$  and ISO<sub>PROG</sub> eliciting favourable duration  $T_{\text{rec}} \geq 38.5^\circ\text{C}$ , duration  $T_{\text{rec}} \geq 39.0^\circ\text{C}$ , change in  $T_{\text{rec}}$  and mean heart rate (HR) when compared to FIXED (Table 2). Isothermic heat acclimation increased the duration spent above the minimum proposed  $T_{\text{rec}}$  of  $38.5^\circ\text{C}$  (Fox et al., 1963) in comparison to FIXED during STHA (duration  $T_{\text{rec}} \geq 38.5^\circ\text{C}$ ; ISO<sub>CONT</sub> = ~32 min session, ISO<sub>PROG</sub> = ~49 min session; FIXED ~24 min session) and LTHA (duration  $T_{\text{rec}} \geq 38.5^\circ\text{C}$ ; ISO<sub>CONT</sub> = ~32 min session, ISO<sub>PROG</sub> = ~46 min session, FIXED ~18 min session), with potential for more complete phenotypic adaptation as a result of consistently longer durations at higher core temperatures (Patterson et al., 2004; Regan et al., 1996; Taylor and Cotter, 2006). Contrary to our hypothesis however, the rate or magnitude of adaptation was not different utilising our between methods, low statistical power was observed for the interaction effect in our data (change HR  $\eta^2 = 0.18$ , sweat loss  $\eta^2 = 0.17$ ,  $T_{\text{sk}}$   $\eta^2 = 0.16$ , all other variables  $\eta^2 \leq 0.10$ ), this may suggest the present study is under powered, or more likely that no difference will be observed when using sample sizes representative of other research in the field, and based upon a priori calculations. The between method statistical analysis implemented may additionally have yielded different observations than that of a repeated measures within method design, with the latter potentially influencing the ability to determine differences in adaptation between STHA and LTHA. FIXED heat acclimation remains a simple method for eliciting adaptation though consistent workloads, however core temperature increases could only be sustained throughout acclimation should a higher exercise intensity, longer exercise duration or elevated exogenous environmental temperature be progressively implemented to counteract the reduced endogenous strain (Galloway and Maughan, 1997; Gibson et al., 2014; Nielsen et al., 1993; Périard et al., 2012). This observation is further evidenced by decreased mean  $T_{\text{rec}}$  and mean  $T_{\text{rec final 60}}$  as adaptation occurred from STHA to LTHA sessions in FIXED.

Within the isothermic methods, failure for ISO<sub>PROG</sub> to confer greater adaptation than ISO<sub>CONT</sub>, suggests a minimum thermoregulatory strain sufficient to elicit physiological adaptations are surpassed by both isothermic, and also fixed methods. Comparison of method administration data (Table 2) suggests individual variability still occurs within methods, particularly isothermic

methods as during STHA, when both ISO<sub>CONT</sub> and ISO<sub>PROG</sub> are performing the same intended protocol, differences are observed in the exercising duration and duration  $T_{\text{rec}} = 38.5^\circ\text{C}$ . This is likely to be due to subtle differences evoked by prescribing workloads based upon a %  $\dot{V}_{\text{O}_{2\text{peak}}}$ . Irrespective of the variation between ISO<sub>CONT</sub> and ISO<sub>PROG</sub>, both elicit greater potentiating stimuli for adaptation than FIXED. Isothermic methods attain the optimal internal temperature for adaptation ( $38.5^\circ\text{C}$ ) for greater durations throughout STHA and LTHA, in line with seminal work in the field (Fox et al., 1963). Isothermic data are similar to that observed during short duration, high exercise intensity heat acclimation ( $75\% \dot{V}_{\text{O}_{2\text{peak}}}$  for 30–35 min day<sup>-1</sup>), which was found to elicit identical adaptation to a longer duration, low exercise intensity heat acclimation method ( $50\% \dot{V}_{\text{O}_{2\text{peak}}}$  for 60 min day<sup>-1</sup>) similar to FIXED (Houmard et al., 1990). Short duration, moderate intensity exercise-heat stress followed by passive rest (isothermic methods) more closely representing competition or training, is equally as effective as longer, lower intensity exposures at inducing adaptations. The benefit of isothermic methods being higher initial workloads reduce exercising durations in comparison to lower intensity continuous fixed methods. The differences in core temperature, and subsequent duration and intensity of work performed between continuous and progressive isothermic methods occur as a result of recent observations that absolute  $\dot{V}_{\text{O}_2}$  is most closely related to metabolic heat production (Smoljanic et al., 2014) (Table 1), and that to ensure equal comparison between groups, workload could be more closely controlled using a workload prescription method other than % (Cramer and Jay, 2014), the authors propose that prescribing heat acclimation utilising workloads known to elicit desired rates of metabolic heat production may reduce variations in heat  $\dot{V}_{\text{O}_{2\text{peak}}}$  gain particularly in an unacclimated individual with relatively lower sweat and evaporative losses. Additionally intermittent exercise, which the latter stages of isothermic protocols can mimic, is known to elicit greater thermal and cardiovascular strain than continuous exercise of the same average intensity (Taylor and Cotter, 2006).

#### 4.3. Physiological mechanisms of heat acclimation

Resting  $T_{\text{rec}}$  and HR reductions were observed towards the latter end of the sessional heat acclimation data (Fig. 2) and during HSTs following STHA and LTHA (Fig. 3). The same magnitude and rate of adaptation indicated that these primary physiological adaptations to heat acclimation regimes occurred using all methods (Garrett et al., 2011; Sawka et al., 2011; Taylor, 2014). Resting and mean exercising  $T_{\text{rec}}$  reduced after STHA, but were not further enhanced after LTHA for all methods in accordance with previous work (Buono et al., 1998; Kampmann et al., 2008). Though comparable with some previous data to determine temperature responses to heat stress (Druyan et al., 2013; Moran et al., 2006), the mean/change  $T_{\text{rec}}$  observed during the HSTs were not as high as observed using alternative protocols to determine physiological responses to heat stress (Magalhães et al., 2010a; Périard et al., 2012). This may have affected the ability of the test to determine core temperature differences augmented by the different heat acclimation methods, or between STHA and LTHA. Additionally, a cycling rather than running heat stress test may have yielded different  $T_{\text{rec}}$  responses specific to the exercise modality of the heat acclimation methods. This may also be true of a test implementing workloads specific to one particular exercise domain or prescribing an intensity more closely reflecting athletic competition or occupational activity. Modified temperature thresholds and plasticity of the hypothalamic neurons within the thermoregulatory centre (Boulant, 2006) and afferent peripheral-central drive (Horowitz, 2014) are proposed mechanisms for this



**Fig. 2.** Mean  $\pm$  SD sessional resting  $T_{\text{rec}}$ , resting HR and sweat loss adaptations to FIXED, ISOCONT, and ISOPROG HA methods. \* Denotes difference from session one ( $p < 0.05$ ).

phenomenon, although the molecular role of prostaglandin E2 (PGE2), cyclooxygenase (COX)-2 and orexin cannot be excluded (Shin et al., 2013). Change in  $T_{\text{rec}}$  did not attenuate from HST1 following STHA or LTHA, therefore adaptations did not offset rate of heat gain (Schlader et al., 2011a). Reduced HR following STHA, combined with lower  $T_{\text{rec}}$ , indicated lower overall physiological strain during HST2, but a further five days of any heat acclimation method did not elicit further adaptations at HST3 (Kampmann et al., 2008).

Sweat losses increased (Fig. 3), likely contributing, alongside reduced skin blood flow (Kenefick et al., 2007) to the reduced mean and peak  $T_{\text{sk}}$  following STHA, but was not enhanced by a LTHA period of 10 days. Though commonly reported as an adaptation following LTHA (Buono et al., 2009; Chinevere et al., 2008), improved sweat loss after STHA is not unique to our data (Machado-Moreira et al., 2006). Our data show enhanced sudomotor function from STHA in all methods of heat acclimation; as such different work duration or intensity does not induce different sweat rate adaptation. It is conceivable that other


thermoregulatory adaptations inhibited the requirement for elevated sweat loss during HST3 rather than a plateau in adaptation being apparent. This is somewhat supported by our daily heat acclimation session data (Fig. 2), whereby increased sweat losses were observed beyond the STHA timescale. It is likely that a lowered internal temperature threshold for sweating was induced by each heat acclimation method (Armstrong and Kenney, 1993; Cotter et al., 1997; Gonzalez et al., 1974; Hessemer et al., 1986; Nadel et al., 1974; Patterson et al., 2004; Roberts et al., 1977; Shido et al., 1999). In conjunction with reduced  $T_{\text{rec}}$ , this adaptation afforded participants an improved centrally-mediated tolerance to exercise-heat stress with the increased sweat loss a consequence of a greater duration spent sweating within the session. This adaptation is facilitated by an earlier onset of sweating (Shido et al., 1999). Heat acclimation is also known to induce peripheral changes at the sweat gland to sweat response during exercise-heat stress (Buono et al., 2009; Fox et al., 1964; Lorenzo and Minson, 2010). Increased cholinergic sensitivity of the eccrine sweat gland or increased glandular hypertrophy is induced by heat acclimation (Sato and Sato, 1983). Sweat adaptations through central (threshold for sweat onset) and peripheral (sweat gland function) mechanisms combine with reductions in the core temperature threshold for cutaneous vasodilation (Buono et al., 1998; Fujii et al., 2012; Hessemer et al., 1986; Nielsen et al., 1997; Yamazaki and Hamasaki, 2003) to confer adaptation decreasing mean and peak  $T_{\text{sk}}$  by sweat evaporation (Fig. 3). These adaptations combined to reduce  $T_{\text{sk}}$ , permitting greater direction of cardiac output to active muscles, as opposed to cutaneous anatomy (González-Alonso et al., 1999) reducing cardiovascular strain in the heat as evidenced by our heart rate data.

The observation that thermal sensation further reduced from STHA to LTHA is potentially beneficial for performance in the heat (Fig. 4), it is believed that thermal discomfort drives true behavioural thermoregulation (Flouris, 2011). Initiation of each of these response pathways during exercise elicit behavioural responses are known to lower work rate (Tucker et al., 2006, 2004). This is an undesirable with regards to optimal performance in the heat however the role of thermal comfort/sensation and pacing are yet to be fully elucidated with contrasting data at present likely due to difference experimental design (Barwood et al., 2014, 2012; Schlader et al., 2011b).

#### 4.4. Applications for practitioners

Environmental conditions in this study were at the upper range of that typically prescribed for heat acclimation. Current practice dictates practitioners aiming to induce heat acclimation would typically administer exercise-heat stress environments at lower ambient temperature and humidity. We propose that in environments of lower heat stress, that isothermal heat acclimation would provide optimal conditions for adaptation compared to fixed intensity methods, where adaptations would not be augmented to the same extent on a sessional basis due to reductions in strain during the acclimation process. This is most relevant during LTHA. With reference to isothermic modes of heat acclimation, as exogenous heat stress reduces, the exercise or training load would increase, potentially reducing the efficacy of this method. Hotter or more humid conditions would offset this, eliciting more rapid increases in  $T_{\text{rec}}$  than cooler conditions (Gibson et al., 2014). Integration of heat acclimation into an athletic taper (Garrett et al., 2011; Mujika et al., 2004) is potentially problematic, requiring acknowledgement of increased work demands of exercise in increasing thermal environments (Galloway and Maughan, 1997), which subsequently decreases with attainment of heat acclimation (Sawka et al., 1983). We suggest practitioners wishing to induce heat acclimation at a time of athletic taper should prescribe





**Fig. 3.** Mean  $\pm$  SD physiological heat stress test data at baseline (HST1), post-STHA sessions (HST2) and post-LTHA (HST3) of FIXED, ISOCONT and ISOPROG. \* Denotes HST significantly different ( $p < 0.05$ ) from HST1 overall.

isothermic heat acclimation under conditions of greater exogenous heat stress than forthcoming competition, to facilitate maximal thermal adaptation for reduced exercise training requirement. The reduction in training volume being an essential component of the taper (Spilsbury et al., 2014), establishing cardiorespiratory, vascular, haematological and neuromuscular changes which ultimately contribute towards optimal performance (Mujika et al., 2004). Additionally due to the greater absolute  $\dot{V}_{O_2}$ , and consequently metabolic heat production, for the same relative workload athletes typically exhibit, the rate of rectal temperature increase is likely to be greater in a trained vs untrained population, reducing the duration taken to reach the 38.5 °C core temperature target facilitating greater rest periods (Table 2).

#### 4.5. Future research directions


Future work could involve the implementation of an isothermic method where workload is implemented using a fixed relative metabolic heat production or relative power, as opposed to a relative workload such as %  $\dot{V}_{O_{2\max}}$ , which may further optimise adaptation by reducing our observed individual variability associated with metabolic heat production and retention (Cramer and Jay, 2014). Additionally due to the linear relationship between HR and  $\dot{V}_{O_2}$ , this physiological measurement may be viable for prescribing work rate during heat acclimation. A comparison of the inducibility of changes in sweat composition, skin blood flow and plasma volume expansion in response to the different heat acclimation modes are yet to be elucidated. It also remains uncertain

whether isothermic heat acclimation is a more efficient method than fixed intensity heat acclimation for preparing highly trained individuals for exercise heat stress (Garrett et al., 2012). Highly trained individuals are likely to be able to sustain the greater absolute workloads required of the isothermic methods, with higher metabolic heat production elevating the rate of core temperature more rapidly (Cramer et al., 2012; Garrett et al., 2012, 2011) and thus, giving greater competition specificity to their acclimation, further enhancing the efficacy of isothermic methods for this population.

#### 5. Conclusions

All heat acclimation methods tested in this study were able to induce the heat-adapted phenotype following five days of heat acclimation and therefore have merit towards attenuating increased physiological strain when exercising in the heat. Based upon our data, the implementation of ten days of heat acclimation did not elicit greater adaptation than five days with the exception of thermal sensation. We have identified that no difference in the extent of adaptation exists between fixed intensity, continuous isothermic, and progressive isothermic heat acclimation methods. Isothermic methods may be more favourable for athletes aiming to integrate heat acclimation into a pre competition taper due to reduced exercise durations and mean session intensities.





**Fig. 4.** Mean  $\pm$  SD perceptual heat stress test data at baseline (HST1), post-STHA sessions (HST2) and post-LTHA (HST3) of FIXED, ISO<sub>CONT</sub> and ISO<sub>PROG</sub>. \* Denotes HST significantly different ( $p < 0.05$ ) from HST1 overall. # Denotes HST significantly different ( $p < 0.05$ ) from HST2 overall.

## Conflict of interests

The authors declare that they have no competing interests.

## Acknowledgements

The authors would like to thank the volunteers for their participation in this investigation.

## References

- Amorim, F., Yamada, P., Robergs, R., Schneider, S., Moseley, P., 2011. Effects of whole-body heat acclimation on cell injury and cytokine responses in peripheral blood mononuclear cells. *Eur. J. Appl. Physiol.* 111, 1609–1618. <http://dx.doi.org/10.1007/s00421-010-1780-4>.
- Aoyagi, Y., McLellan, T.M., Shephard, R.J., 1997. Interactions of physical training and heat acclimation. *The thermophysiology of exercising in a hot climate. Sports Med.* 23, 173–210.
- Armstrong, C.G., Kenney, W.L., 1993. Effects of age and acclimation on responses to passive heat exposure. *J. Appl. Physiol.* 75, 2162–2167.
- Armstrong, L.E., Maresh, C.M., 1991. The induction and decay of heat acclimatization in trained athletes. *Sports Med.* 12, 302–312.
- Barwood, M.J., Corbett, J., White, D., James, J., 2012. Early change in thermal perception is not a driver of anticipatory exercise pacing in the heat. *Br. J. Sports Med.* 46, 936–942. <http://dx.doi.org/10.1136/bjsports-2011-090536>.
- Barwood, M.J., Corbett, J., White, D.K., 2014. Spraying with 0.20% L-menthol does not enhance 5 km running performance in the heat in untrained runners. *J. Sports Med. Phys. Fitness* 54, 595–604.
- Borg, G., Ljunggren, G., Ceci, R., 1985. The increase of perceived exertion, aches and pain in the legs, heart rate and blood lactate during exercise on a bicycle ergometer. *Eur. J. Appl. Physiol. Occup. Physiol.* 54, 343–349. <http://dx.doi.org/10.1007/BF02337176>.

- Boulant, J.A., 2006. Neuronal basis of Hammel's model for set-point thermo-regulation. *J. Appl. Physiol.* 100, 1347–1354. <http://dx.doi.org/10.1152/japplphysiol.01064.2005>.
- Buono, M.J., Heaney, J.H., Canine, K.M., 1998. Acclimation to humid heat lowers resting core temperature. *Am. J. Physiol.* 274, R1295–R1299.
- Buono, M.J., Numan, T.R., Claros, R.M., Brodine, S.K., Kolkhorst, F.W., 2009. Is active sweating during heat acclimation required for improvements in peripheral sweat gland function? *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 297, R1082–R1085. <http://dx.doi.org/10.1152/ajpregu.00253.2009>.
- Burk, A., Timpmann, S., Kreegipuu, K., Tamm, M., Unt, E., Ööpik, V., 2012. Effects of heat acclimation on endurance capacity and prolactin response to exercise in the heat. *Eur. J. Appl. Physiol.*, 4091–4101. <http://dx.doi.org/10.1007/s00421-012-2371-3>.
- Castle, P., Mackenzie, R.W., Maxwell, N., Webborn, A.D.J., Watt, P.W., 2011. Heat acclimation improves intermittent sprinting in the heat but additional pre-cooling offers no further ergogenic effect. *J. Sports Sci.* 29, 1125–1134. <http://dx.doi.org/10.1080/02640414.2011.583673>.
- Castle, P.C., Kularatne, B.P., Brewer, J., Mauger, A.R., Austen, R.A., Tuttle, J.A., Sculthorpe, N., Mackenzie, R.W., Maxwell, N.S., Webborn, A.D.J., 2012. Partial heat acclimation of athletes with spinal cord lesion. *Eur. J. Appl. Physiol.*, 109–115. <http://dx.doi.org/10.1007/s00421-012-2417-6>.
- Chen, T.-I., Tsai, P.-H., Lin, J.-H., Lee, N.-Y., Liang, M.T., 2013. Effect of short-term heat acclimation on endurance time and skin blood flow in trained athletes. *Open access. J. Sports Med.* 4, 161–170. <http://dx.doi.org/10.2147/OAJSM.S45024>.
- Cheung, S.S., McLellan, T.M., 1998. Heat acclimation, aerobic fitness, and hydration effects on tolerance during uncompensable heat stress. *J. Appl. Physiol.* 84, 1731–1739.
- Chinevere, T.D., Kenefick, R.W., Chevront, S.N., Lukaski, H.C., Sawka, M.N., 2008. Effect of heat acclimation on sweat minerals. *Med. Sci. Sports Exerc.* 40, 886–891. <http://dx.doi.org/10.1249/MSS.0b013e3181641c04>.
- Cotter, J.D., Patterson, M.J., Taylor, N.A., 1997. Sweat distribution before and after repeated heat exposure. *Eur. J. Appl. Physiol. Occup. Physiol.* 76, 181–186. <http://dx.doi.org/10.1007/s004210050232>.
- Cramer, M.N., Bain, A.R., Jay, O., 2012. Local sweating on the forehead, but not forearm, is influenced by aerobic fitness independently of heat balance requirements during exercise. *Exp. Physiol.* 97, 572–582. <http://dx.doi.org/10.1113/expphysiol.2011.061374>.
- Cramer, M.N., Jay, O., 2014. Selecting the correct exercise intensity for unbiased comparisons of thermoregulatory responses between groups of different mass and surface area. *J. Appl. Physiol.* 116, 1123–1132. <http://dx.doi.org/10.1152/japplphysiol.01312.2013>.
- Daanen, H.A.M., Jonkman, A.G., Layden, J.D., Linnane, D.M., Weller, A.S., 2011. Optimising the acquisition and retention of heat acclimation. *Int. J. Sports Med.* 32, 822–828. <http://dx.doi.org/10.1055/s-0031-1279767>.
- Dion, T., Savoie, F.A., Asselin, A., Garipey, C., Goulet, E.D.B., 2013. Half-marathon running performance is not improved by a rate of fluid intake above that dictated by thirst sensation in trained distance runners. *Eur. J. Appl. Physiol.* 113, 3011–3020. <http://dx.doi.org/10.1007/s00421-013-2730-8>.
- Drust, B., Waterhouse, J., Atkinson, G., Edwards, B., Reilly, T., 2005. Circadian rhythms in sports performance—an update. *Chronobiol. Int.* 22, 21–44.
- Druyan, A., Ketko, I., Yanovich, R., Epstein, Y., Heled, Y., 2013. Refining the distinction between heat tolerant and intolerant individuals during a heat tolerance test. *J. Therm. Biol.* 38, 539–542. <http://dx.doi.org/10.1016/j.jtherbio.2013.09.005>.
- Du Bois, D., Du Bois, E.F., 1916. A formula to estimate the approximate surface area if height and weight be known. *Arch. Intern. Med.* 17, 863–871.
- Durnin, J.V., Womersley, J., 1974. Body fat assessed from total body density and its estimation from skinfold thickness: measurements on 481 men and women aged from 16 to 72 years. *Br. J. Nutr.* 32, 77–97.
- Flouris, A.D., 2011. Functional architecture of behavioural thermoregulation. *Eur. J. Appl. Physiol.* 111, 1–8. <http://dx.doi.org/10.1007/s00421-010-1602-8>.
- Fox, R.H., Goldsmith, R., Hampton, I.F., Lewis, H.E., 1964. The nature of the increase in sweating capacity produced by heat acclimatization. *J. Physiol.* 171, 368–376.
- Fox, R.H., Goldsmith, R., Kidd, D.J., Lewis, H.E., 1963. Acclimatization to heat in man by controlled elevation of body temperature. *J. Physiol.* 166, 530–547.
- Frank, A., Belokopytov, M., Moran, D., Shapiro, Y., Epstein, Y., 2001. Changes in heart rate variability following acclimation to heat. *J. Basic Clin. Physiol. Pharmacol.* 12, 19–32.
- Fujii, N., Honda, Y., Ogawa, T., Tsuji, B., Kondo, N., Koga, S., Nishiyasu, T., 2012. Short-term exercise-heat acclimation enhances skin vasodilation but not hyperthermic hyperpnea in humans exercising in a hot environment. *Eur. J. Appl. Physiol.* 112, 295–307. <http://dx.doi.org/10.1007/s00421-011-1980-6>.
- Galloway, S.D.R., Maughan, R.J., 1997. Effects of ambient temperature on the capacity to perform prolonged cycle exercise in man. *Med. Sci. Sports Exerc.* 29, 1240–1249. <http://dx.doi.org/10.1097/00005768-199709000-00018>.
- Garrett, A.T., Creasy, R., Rehrer, N.J., Patterson, M.J., Cotter, J.D., 2012. Effectiveness of short-term heat acclimation for highly trained athletes. *Eur. J. Appl. Physiol.* 112, 1827–1837. <http://dx.doi.org/10.1007/s00421-011-2153-3>.
- Garrett, A.T., Goossens, N.G., Rehrer, N.J., Patterson, M.J., Harrison, J., Sammut, I., Cotter, J.D., 2014. Short-term heat acclimation is effective and may be enhanced rather than impaired by dehydration. *Am. J. Hum. Biol.* 26, 311–320. <http://dx.doi.org/10.1002/ajhb.22509>.
- Garrett, A.T., Goossens, N.G., Rehrer, N.J., Rehrer, N.G., Patterson, M.J., Cotter, J.D., 2009. Induction and decay of short-term heat acclimation. *Eur. J. Appl. Physiol.* 107, 659–670. <http://dx.doi.org/10.1007/s00421-009-1182-7>.
- Garrett, A.T., Rehrer, N.J., Patterson, M.J., 2011. Induction and decay of short-term

- heat acclimation in moderately and highly trained athletes. *Sports Med.* 41, 757–771. <http://dx.doi.org/10.2165/11587320-000000000-00000>.
- Gibson, O.R., Dennis, A., Parfitt, T., Taylor, L., Watt, P.W., Maxwell, N.S., 2014. Extracellular Hsp72 concentration relates to a minimum endogenous criteria during acute exercise-heat exposure. *Cell Stress Chaperones* 19, 389–400. <http://dx.doi.org/10.1007/s12192-013-0468-1>.
- Gonzalez, R.R., Pandolf, K.B., Gage, A.P., 1974. Heat acclimation and decline in sweating during humidity transients. *J. Appl. Physiol.* 36, 419–425.
- González-Alonso, J., Teller, C., Andersen, S.L., Jensen, F.B., Hyldig, T., Nielsen, B., 1999. Influence of body temperature on the development of fatigue during prolonged exercise in the heat. *J. Appl. Physiol.* 86, 1032–1039.
- Hessemer, V., Zeh, A., Brück, K., 1986. Effects of passive heat adaptation and moderate sweatless conditioning on responses to cold and heat. *Eur. J. Appl. Physiol. Occup. Physiol.* 55, 281–289. <http://dx.doi.org/10.1007/BF02343800>.
- Hom, L.L., Lee, E.C.-H., Apicella, J.M., Wallace, S.D., Emmanuel, H., Klau, J.F., Poh, P.Y.S., Marzano, S., Armstrong, L.E., Casa, D.J., Maresh, C.M., 2012. Eleven days of moderate exercise and heat exposure induces acclimation without significant HSP70 and apoptosis responses of lymphocytes in college-aged males. *Cell Stress Chaperones* 17, 29–39. <http://dx.doi.org/10.1007/s12192-011-0283-5>.
- Horowitz, M., 2014. Heat acclimation, epigenetics, and cytoprotection memory. *Compr. Physiol.* 4, 199–230. <http://dx.doi.org/10.1002/cphy.c130025>.
- Houmard, J.A., Costill, D.L., Davis, J.A., Mitchell, J.B., Pascoe, D.D., Robergs, R.A., 1990. The influence of exercise intensity on heat acclimation in trained subjects. *Med. Sci. Sports Exerc.* 22, 615–620.
- Kampmann, B., Bröde, P., Schütte, M., Griefahn, B., 2008. Lowering of resting core temperature during acclimation is influenced by exercise stimulus. *Eur. J. Appl. Physiol.* 104, 321–327. <http://dx.doi.org/10.1007/s00421-007-0658-6>.
- Kenefick, R.V., Chevront, S.N., Sawka, M.N., 2007. Thermoregulatory function during the marathon. *Sports Med.* 37, 312–315.
- Kresfelder, T.L., Claassen, N., Cronjé, M.J., 2006. Hsp70 Induction and hsp70 gene polymorphisms as indicators of acclimatization under hyperthermic conditions. *J. Therm. Biol.* 31, 406–415. <http://dx.doi.org/10.1016/j.jtherbio.2006.02.001>.
- Lorenzo, S., Halliwill, J.R., Sawka, M.N., Minson, C.T., 2010. Heat acclimation improves exercise performance. *J. Appl. Physiol.* 109, 1140–1147. <http://dx.doi.org/10.1152/jappphysiol.00495.2010>.
- Lorenzo, S., Minson, C.T., 2010. Heat acclimation improves cutaneous vascular function and sweating in trained cyclists. *J. Appl. Physiol.* 109, 1736–1743. <http://dx.doi.org/10.1152/jappphysiol.00725.2010>.
- Machado-Moreira, C.A., Vimieiro-Gomes, A.C., Silami-Garcia, E., Lima, N.R.V., Rodrigues, L.O.C., 2006. Possible biphasic sweating response during short-term heat acclimation protocol for tropical natives. *J. Physiol. Anthropol.* 25, 215–219. <http://dx.doi.org/10.2114/jpa.2.25.215>.
- Magalhães, F.D.C., Amorim, F.T., Passos, R.L.F., Fonseca, M.A., Oliveira, K.P.M., Lima, M.R.M., Guimarães, J.B., Ferreira-Júnior, J.B., Martini, A.R.P., Lima, N.R.V., Soares, D.D., Oliveira, E.M., Rodrigues, L.O.C., 2010a. Heat and exercise acclimation increases intracellular levels of Hsp72 and inhibits exercise-induced increase in intracellular and plasma Hsp72 in humans. *Cell Stress Chaperones* 15, 885–895. <http://dx.doi.org/10.1007/s12192-010-0197-7>.
- Magalhães, F.D.C., Passos, R.L.F., Fonseca, M.A., Oliveira, K.P.M., Ferreira-Júnior, J.B., Martini, A.R.P., Lima, M.R.M., Guimarães, J.B., Baraúna, V.G., Silami-Garcia, E., Rodrigues, L.O.C., 2010b. Thermoregulatory efficiency is increased after heat acclimation in tropical natives. *J. Physiol. Anthropol.* 29, 1–12.
- Marshall, H.C., Campbell, S.A., Roberts, C.W., Nimmo, M.A., 2007. Human physiological and heat shock protein 72 adaptations during the initial phase of humid-heat acclimation. *J. Therm. Biol.* 32, 341–348. <http://dx.doi.org/10.1016/j.jtherbio.2007.04.003>.
- Martinez, R., Jones, D., Hodge, D., Buono, M.J., 2012. Blocking the beta-adrenergic system does not affect sweat gland function during heat acclimation. *Auton. Neurosci.* 169, 113–115. <http://dx.doi.org/10.1016/j.autneu.2012.05.007>.
- Moran, D.S., Eli-Berchoer, L., Heled, Y., Mendel, L., Schocina, M., Horowitz, M., 2006. Heat intolerance: does gene transcription contribute? *J. Appl. Physiol.* 100, 1370–1376. <http://dx.doi.org/10.1152/jappphysiol.01261.2005>.
- Mujika, I., Padilla, S., Pyne, D., Busso, T., 2004. Physiological changes associated with the pre-event taper in athletes. *Sports Med.* 34, 891–927 (doi:34133 [pii]).
- Nadel, E.R., Pandolf, K.B., Roberts, M.F., Stolwijk, J.A., 1974. Mechanisms of thermal acclimation to exercise and heat. *J. Appl. Physiol.* 37, 515–520.
- Nielsen, B., Hales, J.R., Strange, S., Christensen, N.J., Warberg, J., Saltin, B., 1993. Human circulatory and thermoregulatory adaptations with heat acclimation and exercise in a hot, dry environment. *J. Physiol.* 460, 467–485.
- Nielsen, B., Strange, S., Christensen, N.J., Warberg, J., Saltin, B., 1997. Acute and adaptive responses in humans to exercise in a warm, humid environment. *Pflügers Arch.* 434, 49–56.
- Pandolf, K.B., 1979. Effects of physical training and cardiorespiratory physical fitness on exercise-heat tolerance: recent observations. *Med. Sci. Sports* 11, 60–65.
- Patterson, M.J., Stocks, J.M., Taylor, N.A.S., 2014. Whole-body fluid distribution in humans during dehydration and recovery, before and after humid-heat acclimation induced using controlled hyperthermia. *Acta Physiol. (Oxf.)* 210, 899–912. <http://dx.doi.org/10.1111/apha.12214>.
- Patterson, M.J., Stocks, J.M., Taylor, N.A.S., 2004. Humid heat acclimation does not elicit a preferential sweat redistribution toward the limbs. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 286, R512–R518. <http://dx.doi.org/10.1152/ajpregu.00359.2003>.
- Périard, J.D., Ruell, P., Caillaud, C., Thompson, M.W., 2012. Plasma Hsp72 (HSPA1A) and Hsp27 (HSPB1) expression under heat stress: influence of exercise intensity. *Cell Stress Chaperones* 17, 375–383. <http://dx.doi.org/10.1007/s12192-011-0313-3>.
- Ramanathan, N.L., 1964. A new weighting system for mean surface temperature of the human body. *J. Appl. Physiol.* 19, 531–533.
- Regan, J.M., Macfarlane, D.J., Taylor, N.A., 1996. An evaluation of the role of skin temperature during heat adaptation. *Acta Physiol. Scand.* 158, 365–375. <http://dx.doi.org/10.1046/j.1365-201X.1996.561311000.x>.
- Roberts, M.F., Wenger, C.B., Stolwijk, J.A., Nadel, E.R., 1977. Skin blood flow and sweating changes following exercise training and heat acclimation. *J. Appl. Physiol.* 43, 133–137.
- Sandström, M.E., Siegler, J.C., Lovell, R.J., Madden, L., McNaughton, L., 2008. The effect of 15 consecutive days of heat-exercise acclimation on heat shock protein 70. *Cell Stress Chaperones* 13, 169–175. <http://dx.doi.org/10.1007/s12192-008-0022-8>.
- Sato, K., Sato, F., 1983. Individual variations in structure and function of human eccrine sweat gland. *Am. J. Physiol.* 245, R203–R208.
- Sawka, M.N., Burke, L.M., Eichner, E.R., Maughan, R.J., Montain, S.J., Stachenfeld, N.S., 2007. American College of Sports Medicine position stand. Exercise and fluid replacement. *Med. Sci. Sports Exerc.* 39, 377–390. <http://dx.doi.org/10.1249/mss.0b013e31802ca597>.
- Sawka, M.N., Leon, L.R., Montain, S.J., Sanna, L.A., 2011. Integrated physiological mechanisms of exercise performance, adaptation, and maladaptation to heat stress. *Compr. Physiol.* 1, 1883–1928. <http://dx.doi.org/10.1002/cphy.c100082>.
- Sawka, M.N., Pandolf, K.B., Avellini, B.A., Shapiro, Y., 1983. Does heat acclimation lower the rate of metabolism elicited by muscular exercise? *Aviat. Space Environ. Med.* 54, 27–31.
- Schlader, Z.J., Raman, A., Morton, R.H., Stannard, S.R., Mündel, T., 2011a. Exercise modality modulates body temperature regulation during exercise in uncompensable heat stress. *Eur. J. Appl. Physiol.* 111, 757–766. <http://dx.doi.org/10.1007/s00421-010-1692-3>.
- Schlader, Z.J., Simmons, S.E., Stannard, S.R., Mündel, T., 2011b. The independent roles of temperature and thermal perception in the control of human thermoregulatory behavior. *Physiol. Behav.* 103, 217–224. <http://dx.doi.org/10.1016/j.physbeh.2011.02.002>.
- Shido, O., Sugimoto, N., Tanabe, M., Sakurada, S., 1999. Core temperature and sweating onset in humans acclimated to heat given at a fixed daily time. *Am. J. Physiol.* 276, R1095–R1101.
- Shin, Y.O., Lee, J.B., Min, Y.K., Yang, H.M., 2013. Heat acclimation affects circulating levels of prostaglandin E2, COX-2 and orexin in humans. *Neurosci. Lett.* 542, 17–20. <http://dx.doi.org/10.1016/j.neulet.2013.03.017>.
- Siri, W.E., 1956. The gross composition of the body. *Adv. Biol. Med. Phys.* 4, 239–280.
- Smoljanic, J., Morris, N.B., Dervis, S., Jay, O., 2014. Running economy, not aerobic fitness, independently alters thermoregulatory responses during treadmill running. *J. Appl. Physiol.* 117, 1451–1459. <http://dx.doi.org/10.1152/jappphysiol.00665.2014>.
- Spilsbury, K.L., Fudge, B.W., Ingham, S.A., Faulkner, S.H., Nimmo, M.A., 2014. Tapering strategies in elite British endurance runners. *Eur. J. Sports Sci.* 1–7. <http://dx.doi.org/10.1080/17461391.2014.955128>.
- Taylor, L., Midgley, A.W., Christmas, B., Hilman, A.R., Madden, L.A., Vince, R.V., McNaughton, L.R., 2011. Daily hypoxia increases basal monocyte HSP72 expression in healthy human subjects. *Amino Acids* 40, 393–401. <http://dx.doi.org/10.1007/s00726-010-0644-x>.
- Taylor, N., Cotter, J., 2006. Heat adaptation: guidelines for the optimisation of human performance. *Int. Sports Med. J.* 7, 33–57.
- Taylor, N.A.S., 2014. Human heat adaptation. *Compr. Physiol.* 4, 325–365.
- Toner, M.M., Drolet, L.L., Pandolf, K.B., 1986. Perceptual and physiological responses during exercise in cool and cold water. *Percept. Mot. Skills* 62, 211–220.
- Tucker, R., Marle, T., Lambert, E.V., Noakes, T.D., 2006. The rate of heat storage mediates an anticipatory reduction in exercise intensity during cycling at a fixed rating of perceived exertion. *J. Physiol.* 574, 905–915. <http://dx.doi.org/10.1113/jphysiol.2005.101733>.
- Tucker, R., Rauch, L., Harley, Y.X.R., Noakes, T.D., 2004. Impaired exercise performance in the heat is associated with an anticipatory reduction in skeletal muscle recruitment. *Pflügers Arch.* 448, 422–430. <http://dx.doi.org/10.1007/s00424-004-1267-4>.
- Watkins, A.M., Cheek, D.J., Harvey, A.E., Blair, K.E., Mitchell, J.B., 2008. Heat acclimation and HSP-72 expression in exercising humans. *Int. J. Sports Med.* 29, 269–276.
- Yamada, P.M., Amorim, F.T., Moseley, P., Robergs, R., Schneider, S.M., 2007. Effect of heat acclimation on heat shock protein 72 and interleukin-10 in humans. *J. Appl. Physiol.* 103, 1196–1204. <http://dx.doi.org/10.1152/jappphysiol.00242.2007>.
- Yamazaki, F., Hamasaki, K., 2003. Heat acclimation increases skin vasodilation and sweating but not cardiac baroreflex responses in heat-stressed humans. *J. Appl. Physiol.* 95, 1567–1574. <http://dx.doi.org/10.1152/jappphysiol.00063.2003>.



**Oliver Gibson** began his Ph.D. at the University of Brighton in 2010 after receiving his master's degree in Applied Exercise Physiology in 2009 and undergraduate degree in Sport and Exercise Science in 2007. Oliver's Ph.D. examines the cellular stress response to acute and chronic exercise heat stress. Oliver is a technical instructor for the undergraduate and postgraduate Sport and Exercise Science degrees at the University of Brighton, and combines his thermoregulatory doctoral research with research allied to intermittent sprint performance and training in hypoxia, and ergogenic aids to improve endurance performance.



**Jessica Mee** received her undergraduate degree in Sport Science from the University of Brighton in 2010. Jessica began her Ph.D. at the University of Brighton in 2011, which is examining the thermo tolerance and adaptation to the heat in females. Jessica is a technical instructor for the undergraduate and postgraduate Sport and Exercise Science degrees at the University of Brighton.



**Peter Watt** is a reader at the University of Brighton within the Centre for Sport and Exercise Science and Medicine (SESAME) and is the research theme champion for Exercise and Health. Peter's research primarily focuses on the application of stable isotope methods to measure metabolic and physiological changes occurring in humans during exercise, with application to health related problems, e.g. diabetes, obesity. Peter is also research active in areas including exercise and nutritional interactions which affect muscle growth and function, ammonia and fatigue, the effects of exercise and hypoxia on whole body metabolism in diabetes, and physiological responses to heat stress.



**James Tuttle** is a molecular exercise physiology Ph.D. student and laboratory technician at the University of Bedfordshire. James' Ph.D. investigated the contribution of heat and mechanical stress to heat shock protein (HSP) gene expression and mechanisms through which HSPs attenuate exercise induced muscle damage. James' other research interests include the applied exercise physiology of endurance sports particularly middle and long distance running.



**Neil Maxwell** joined the University of Brighton as a lecturer in sport and exercise science in 1997. Neil continues to lecture undergraduate and postgraduate students, predominantly in the area of exercise and environmental physiology and research methods. Neil is research active and has published extensively in the international, scientific literature in areas allied to thermal and hypoxic stress and how the body tolerates each, particularly during exercise. He is an approved higher degrees supervisor with MPhil/Ph.D. completions, external examination experience and a bank of existing postgraduate research students.



**Lee Taylor** is a senior lecturer at the University of Bedfordshire. Lee is a member of the Institute of Sport and Physical Activity (ISPAR Bedford) and Muscle Cellular and Molecular Physiology (MCMP) Applied Sport and Exercise Physiology (ASEP) research groups. Lee's research interests are allied to the *in vivo* stress response to both environmental (hypoxia, hypothermia, hyperthermia and hyperbaria) and/or exercise stress. Specific "stress response" related interests include heat shock proteins (HSPs), disturbances to redox balance and other pro-inflammatory cytokine responses (IL-6, TNF- $\alpha$ , etc.). Additionally Lee has published data implementing the use of various strategies (pre-cooling,

pharmacological agents, acclimation strategies, etc.) to ameliorate the negative impact such environments can have on exercise performance and occupational pursuits.